

***FOREST BOTANY MANUAL***  
***MODULE 4***  
***FREYCINET REGION***



**2005**

**FPA**  
FOREST PRACTICES AUTHORITY

**FOREST BOTANY MANUAL:  
MODULE 4 – FREYCINET REGION**

**2005**

© Forest Practices Authority, Tasmania

Information in the *Forest Botany Manual* may be copied for personal use or published for educational or training purposes provided that any extracts are fully acknowledged.

The suggested citation for the *Forest Botany Manual* is:

Forest Practices Authority (2005). *Forest Botany Manual*. Forest Practices Authority, Tasmania.

The suggested citation for this module is:

Forest Practices Authority (2005). *Forest Botany Manual: Module 4 – Freycinet Region*. Forest Practices Authority, Tasmania.

---

Published by the Forest Practices Authority, 30 Patrick Street, Hobart – Tasmania – 7000

Phone: (03) 6233 7966; Fax: (03) 6233 7954

Email: [info@fpa.tas.gov.au](mailto:info@fpa.tas.gov.au); Website: [www.fpa.tas.gov.au](http://www.fpa.tas.gov.au)

---

## TABLE OF CONTENTS

<b>INTRODUCTION</b> .....	<b>1</b>
<b>Section 1 OVERVIEW OF FREYCINET REGION</b> .....	<b>2</b>
<b>Section 2 FOREST COMMUNITIES</b> .....	<b>4</b>
Using the keys and tables .....	4
Conservation priorities .....	5
Explanatory notes .....	6
<b>KEY TO VEGETATION TYPES AND FOREST COMMUNITIES</b> .....	<b>10</b>
Key to forest types.....	10
Key to non-forest types.....	10
<b>2.1 RAINFOREST COMMUNITIES</b> .....	<b>12</b>
Key to rainforest communities.....	12
Conservation priorities and attributes of rainforest communities .....	13
<b>2.2 SWAMP FOREST AND RELATED FOREST OR SCRUB COMMUNITIES</b> .....	<b>14</b>
Key to swamp forest and related forest or scrub communities .....	14
Conservation priorities and attributes of swamp forest and related forest or scrub communities .....	15
<b>2.3 OTHER FOREST OR SCRUB COMMUNITIES</b> .....	<b>16</b>
Key to other forest or scrub communities.....	16
Conservation priorities and attributes of other forest or scrub communities .....	17
<b>2.4 MIXED FOREST COMMUNITIES</b> .....	<b>18</b>
Key to mixed forest communities.....	18
Conservation priorities and attributes of mixed forest communities .....	19
<b>2.5 WET SCLEROPHYLL FOREST COMMUNITIES</b> .....	<b>21</b>
Key to wet sclerophyll forest communities .....	21
Conservation priorities and attributes of wet sclerophyll forest communities.....	23
<b>2.6 DRY SCLEROPHYLL FOREST AND WOODLAND COMMUNITIES</b> .....	<b>27</b>
Recognising the different dry sclerophyll understoreys .....	27
Key to dry sclerophyll forest and woodland communities .....	28
Conservation priorities and attributes of dry sclerophyll forest / woodland communities .....	30
<b>Section 3 PRIORITY SPECIES</b> .....	<b>36</b>
Forest quality moderate to high .....	37
Forest quality low to moderate .....	37
Forest quality very low or non-forest .....	38
<b>Section 4 SITES OF POTENTIAL SIGNIFICANCE FOR FLORA</b> .....	<b>42</b>
<b>Section 5 EVALUATION OF OTHER FLORA ISSUES</b> .....	<b>45</b>
Weed and disease management .....	45
Remnant forest and woodland .....	46
Management of forestry operations in vicinity of reserves.....	46
<b>Section 6 ASSESSING THE FLORA VALUES OF AN AREA</b> .....	<b>48</b>

## INTRODUCTION

Tasmania is divided into eight bioregions on the basis of the State's biogeography. Separate Forest Botany Modules have been developed for these regions.

This module deals with the Freycinet Region. It covers subjects relevant to conservation of flora, as required by the Tasmanian *Forest Practices Code* (2000), the *Forest Practices Act* and other legislation and processes.

The module is designed to assist Forest Practices Officers (FPOs), and others involved with forest management, to prepare Forest Practices Plans (FPPs) for sites within the region. The information can also be used for other purposes (e.g. management planning for reserves, preparation of property plans).

The module is divided into six sections, which follow the format of the FPP *Flora Evaluation Sheet*:

**Section 1** gives a brief overview of Freycinet Region.

**Section 2** provides keys to forest and non-forest vegetation, and more detailed keys to forest communities. Tables indicate conservation priorities for forest communities.

**Section 3** lists plant species that have a priority for conservation in the region - most of these are species listed on the Tasmanian *Threatened Species Protection Act 1995*.

**Section 4** indicates sites of potential significance for flora conservation. These are environments that are often associated with species or communities that have a priority for conservation.

**Section 5** discusses some other issues (e.g. weed and disease management) that may need to be considered by FPOs, to ensure that the operation complies with botanical requirements of the *Forest Practices Code* and other policies.

**Section 6** summarises the evaluation process and indicates the steps that need to be taken after a FPO has assessed the FPP area. It also indicates whether specialist advice is required.

The processes used to determine if communities, species and sites of potential significance are present in an area, will also capture those National Estate flora values (as identified in the Tasmanian Regional Forest Agreement) that have the potential to be affected by operations requiring FPPs.

Module 1 of the *Forest Botany Manual* gives background information relevant to users of the regional modules. The Manual is supported by information on the Forest Practices Authority (FPA) website, including a gallery containing images of many threatened species, and species used to identify vegetation types and forest communities. An ongoing series of Flora Technical Notes also covers aspects of vegetation management in Tasmanian forests. The Manual provides links to several external websites – the FPA website will maintain updates to these sites, and should be consulted if there are problems accessing the links given in the Manual.

Queries and comments about the format or content of the *Forest Botany Manual* should be referred to the FPA's Senior Botanist. Queries and notifications about vegetation in operational areas should generally be referred to the Senior Ecologist.

### Contact details for the Forest Practices Authority's Botany and Ecology programs:

**Forest Practices Authority,  
30 Patrick Street,  
Hobart, Tasmania 7000**

**Phone: (03) 62337804 (Botany); (03) 62164454 or 62337870 (Ecology)**

**Fax: (03) 62337954**

**Email: [info@fpa.tas.gov.au](mailto:info@fpa.tas.gov.au) ; Website: [www.fpa.tas.gov.au](http://www.fpa.tas.gov.au)**

## Section 1 OVERVIEW OF FREYCINET REGION

Freycinet Region primarily covers coastal and hinterland areas of southeastern Tasmania. It includes Freycinet, Tasman and Forestier Peninsulas, and associated islands. The Eastern Tiers form a dissected and mountainous backbone to the region, which extends west to the more arable (and agricultural) Midlands. The region contains a wide range of rock types and landforms, and steep climatic gradients at regional and local levels. Consequently, the native vegetation is also diverse. Land use and fire history have also influenced the extent, structure and composition of the vegetation.


Dry sclerophyll forests and woodlands are the most widespread vegetation type in Freycinet Region. Wet sclerophyll forests are common in moister upland areas and on Forestier and Tasman Peninsulas, and also occupy shaded environments in drier parts of the region. Mixed forests (eucalypt forest with a rainforest understorey) and rainforests are very restricted in distribution, and are largely confined to fire-shadow areas in more humid parts of the region. Small areas of other forest types (e.g. swamp forest, Oyster Bay pine forest and she-oak forest) occur confined to localised habitats. Non-forest vegetation occurring in Freycinet Region includes native grassland (mainly dry inland areas), heath, scrub and moorland (mainly infertile coastal environments), salt marshes and wetlands.

Freycinet Region contains about a thousand species of vascular plants that are native to Tasmania. About 150 of these species are Tasmanian endemics. A high proportion of the endemic species are associated with dry or exposed sites on dolerite, a rock type virtually absent from the southeastern Australian mainland. Several endemic species, and some non-endemic species, have their only recorded distribution in the Freycinet Region. The non-vascular flora of Freycinet Region (i.e. mosses, liverworts and lichens) is poorly known compared to the vascular flora. Non-vascular diversity is highest in rainforest and mixed forest communities.

The native vegetation of more arable environments has been most affected by settlement and agriculture. The least agriculturally viable areas, such as coastal landforms and the extensive dolerite uplands, support large areas of native forest. Forestry is a significant land use in many of the remaining wetter and drier forests. Plantation establishment has mainly occurred in wetter parts of the region. Extensive grazing is practised in more open forests and woodlands. The root rot pathogen *Phytophthora cinnamomi* is a threat to the biota in parts of the region.

A substantial proportion of the native vegetation in Freycinet Region is formally reserved. Larger reserves containing native forest include Freycinet National Park, Douglas-Apsley NP, Abel Tasman NP, Tooms Lake Forest Reserve, Buxton River FR and Sandspit River FR. Recent additions to the public reserve system have improved the conservation status of many species and communities in the Freycinet Region. Some forest communities and species remain threatened or poorly reserved, and require additional protection through prescription or reservation on public land (e.g. through Special Management Zoning on State forest) and private land (e.g. through agreements developed through the Private Forest Reserves Program).

An overview of the vegetation of Freycinet Region and its relationship with the environment is given in Davies (1988). Some useful references on plant species and vegetation types are given in Module 1 and *Flora Technical Note 2*.



**Location of Freycinet Region**

The eastern and southern boundaries are formed by the coastline and Pittwater Estuary to Sorell. The western boundary is initially formed by the Tasman Highway to Woodsdale Road, which is followed north to the Little Swanport River. The river is followed downstream to Green Tier Creek, which is followed north to Tooms Lake Road. The road is followed west to the 300 m contour, which is followed north for 80 km, to an unnamed creek southwest of Morgans Sugarloaf. This creek is followed north to the South Esk River, which is followed west for 1 km to Horse Paddock Gully. The western tributary of this gully is followed to the 300 m contour. The contour forms the northern boundary of the region, trending northeast along the Fingal Valley (about 120 km), until it reaches Margisons Creek near St Marys. Margisons Creek is followed upstream to a small saddle (south of St Patricks Head) - this is crossed to Banticks Creek, which is followed to the coast.

Numbers refer to 1:25000 Tasmaps.

## Section 2 FOREST COMMUNITIES

This section provides keys to the native vegetation types and native forest communities occurring within Freycinet Region. Tables indicate the conservation status of forest communities. The explanatory notes should be read before the keys and tables are used. The Forest Practices Plan *Flora Evaluation Sheet* indicates when FPOs need specialist botanical advice, when communities that may have a priority for conservation could be affected by forestry operations.

The keys are based on species and other vegetation characteristics that should be familiar to FPOs and field workers. Illustrations of species used in the keys are given in several publications listed in *Flora Technical Note 2*. The FPA website also carries scanned images of diagnostic species. The common and scientific names of all species used in the keys are listed in Appendix 6 of Module 1.

This section does not cover existing plantations (hardwood or softwood) or areas of exotic vegetation (e.g. pasture). Botanical advice relating to communities is not needed if the proposed operation will only affect these vegetation types. However, FPOs need to consider if other botanical values (e.g. threatened species) have the potential to occur on such sites.

Some non-forest vegetation in Freycinet Region has a high priority for conservation, contains threatened species or is very susceptible to disturbance or disease. There are guidelines in the *Forest Practices Code* to avoid disturbance to localised environments that contain these vegetation types. Seek botanical advice in all cases where native non-forest vegetation will be affected by forestry operations.

### USING THE KEYS AND TABLES

The forests occurring in Freycinet Region have been divided into several broad forest types:

- Rainforest;
- Swamp forest and related forest or scrub;
- Other forest or scrub;
- Mixed forest (eucalypt forest with rainforest species also prominent);
- Wet sclerophyll forest;
- Dry sclerophyll forest and woodland.

Different researchers have classified each forest type into several communities, on the basis of the composition and structure of the overstorey and understorey. These are called **floristic communities** in the Manual. The floristic communities can be allocated to the forest communities that were described and mapped for the RFA - these are called **RFA communities** in the Manual. There is generally good correlation between floristic communities and RFA communities, but this is not always the case.

**Use the keys to determine:**

- **The forest types and non-forest vegetation types present in the area;**
- **The floristic communities present in each forest type.**

**Use the associated tables to determine:**

- **The RFA communities present (based on the floristic communities identified);**
- **The conservation priority of each of the floristic communities and RFA communities.**

Some forest communities are particularly susceptible to the root rot pathogen *Phytophthora cinnamomi* – these are also identified in the tables (see discussion on page 10).

A typical native forest coupe in Freycinet Region is likely to contain 3 to 5 floristic communities, the number being largely related to variation in the environment (e.g. landform, rock type, disturbance history). There will usually be more floristic communities than RFA communities in any given area, because the RFA communities are less finely differentiated. It is important to identify the floristic communities, as they give a much better picture of the variation in the region's forests than the RFA community classification. For example, in Freycinet Region there are ten mixed forest or wet sclerophyll forest floristic communities dominated by *E. delegatensis*. These are all included in the RFA community "tall *E. delegatensis* forest." Most of the *E. delegatensis* floristic communities are well reserved in the region, but two are poorly reserved.

It is important to recognise that any system of vegetation classification imposes a taxonomy on something that varies continuously in nature. In addition, our knowledge of Tasmania's vegetation is far from complete. Consequently, FPOs will inevitably come across forest vegetation that does not key out easily. There are a few reasons for this. They include:

- the community may not have been previously recorded from Freycinet Region;
- the community may be close to a particular community given in the key, but in the area assessed may lack a species or other characteristic that allows it to be keyed out to that community (this may happen if fire or other disturbance has altered the structure or composition of the vegetation);
- the community may be intermediate between two communities given in the keys (this may happen if vegetation is sampled in transition zones).

**It is essential that proposed operational areas are field assessed to determine the range of forest and non-forest vegetation that they contain.** Using a combination of the RFA vegetation map, PI maps, aerial photographs, geology maps, topographic maps and local information, will give a good indication of where different communities may occur in an FPP area.

Contact the FPA Botanist or FPA Ecologist if you have problems identifying communities, providing details of the vegetation and site. References given for each forest type also contain useful information.

## CONSERVATION PRIORITIES

Conservation priorities for forest communities are based on the requirements and findings of the RFA and associated processes (e.g. identification of communities as Rare, Vulnerable or Endangered) and known distribution of communities in formal reserves (see Module 1).

FPA advice regarding a priority community in an operational area will depend on many factors. They include: conservation status and distribution of the community; the condition of the vegetation; the nature of the proposed operation; presence of other values; and legislative or policy requirements. In some cases, no changes to plans will be needed, in others prescriptions or reservation will be required.

### Conservation priorities for floristic communities

Priority	Explanation	General course of action
<b>A</b>	Community may be inadequately reserved in Tasmania, and/or may have a very high conservation priority in the region.	Seek botanical advice in all cases if an area is thought to contain a Priority A community.
<b>B</b>	Community may be inadequately reserved in the region, but is adequately reserved elsewhere in Tasmania.	Seek botanical advice if an area is thought to contain a Priority B community <u>and</u> the site will not be regenerated to native forest.
<b>Non-priority (np)</b>	Community is adequately reserved in Tasmania and in the region.	Unless priority species (Section 3) or other flora values are thought to be present, there is generally no need to seek botanical advice if an area only contains non-priority communities.

### Conservation priorities for RFA communities

Priority	Explanation	General course of action
<b>Y</b>	The RFA has identified that additional Statewide conservation is required for the community (oldgrowth and non-oldgrowth).	Seek botanical advice in all cases if an area is thought to contain a Priority Y community.
<b>Yog</b>	The RFA has identified that additional Statewide conservation is required for the oldgrowth component of the community.	Seek botanical advice where the community is oldgrowth, or other flora values are thought to be present.
<b>Non-priority (N)</b>	The RFA has not identified that additional Statewide conservation is required for the community.	Unless priority species (Section 3) or other flora values are thought to be present, there is generally no need to seek botanical advice if an area only contains non-priority communities.

RFA processes have identified communities that are Rare (R), Vulnerable (V) or Endangered (E) at a Statewide level. These are identified (\*) in the tables that indicate the conservation priorities and attributes of the different forest types (see column dealing with conservation status of the RFA community). RVE communities in Tasmania are listed in Module 1 (Appendix 3).

The RFA lists several forest communities that require further protection on public land in Tasmania. Most of these communities also have a high priority for conservation on private land.

The table below lists all RVE communities, and other communities that require protection on public land, that have been recorded from Freycinet Region.

- R E *Melaleuca ericifolia* coastal swamp forest
- R E *Notelaea ligustrina* and/or *Pomaderris apetala* closed forest
- E Shrubby *E. ovata* - *E. viminalis* forest
- E *E. viminalis* wet forest
- R V *E. viminalis* and/or *E. globulus* coastal shrubby forest
- V *Callitris rhomboidea* forest
- V *E. brookeriana* wet forest
- V *E. amygdalina* forest on sandstone
- V Inland *E. amygdalina* - *E. viminalis* - *E. pauciflora* forest / woodland on Cainozoic deposits
- V Inland *E. tenuiramis* forest
- V Grassy *E. globulus* forest
- E. viminalis* grassy forest / woodland
- E. rodwayi* forest
- Allocasuarina verticillata* forest
- E. amygdalina* forest on mudstone (oldgrowth only)
- E. pauciflora* forest on dolerite (oldgrowth only)
- E. sieberi* forest on granite (oldgrowth only)
- E. sieberi* forest on other substrates (oldgrowth only)
- E. viminalis* - *E. ovata* - *E. amygdalina* - *E. obliqua* damp sclerophyll forest (oldgrowth only).

The tables in this section of the Freycinet Module indicate that all occurrences of these communities need to be referred to FPA. The other RFA communities that require referral to FPA have been identified through other analyses as having some priority for conservation within the region.

There are constraints on conversion of RVE forest communities (and RVE non-forest communities). There will be restrictions on further conversion of other forest communities if their clearance approaches the limits set by Tasmania's Permanent Forest Estate Policy (monitored by FPA).

## EXPLANATORY NOTES

Some additional notes that will help FPOs to assess areas and use the keys and tables are given below.

### Sources of information

There are many sources of information to indicate which vegetation types and forest communities occur in a FPP area. Assessments and surveys conducted prior to preparing FPPs will generally provide enough information to identify the communities. Published and unpublished reports and botanical data from various databases may also be useful. Distribution notes given in the tables may help confirm community identifications.

Broad scale vegetation maps such as the RFA Forest Communities Map and TASVEG maps are available through DPIWE GTSpot database and Forestry Tasmania's NewCONSERVE database. Details for accessing these databases are given in Module 1. These maps may give a useful indication of the vegetation in a FPP area, but the scale of mapping means that they are often inaccurate at the coupe level. They rarely pick up localised occurrences of communities (which may have high conservation significance), and they will not allow floristic communities to be identified.

Further information about the different forest types is provided in the major references cited in the text. *Flora Technical Note 2* provides other references on forest and non-forest vegetation.

## How big is a forest community?

In preparing FPPs, the **minimum** area of forest that should be identified as a distinct community is **1 ha** (this includes contiguous areas of the community that extend beyond the FPP boundary). However, botanical advice should be sought for smaller areas of non-forest vegetation (e.g. *Sphagnum* peatlands).

Small areas of communities can be easily missed during surveys of FPP areas, though the chances of this are reduced by good sampling across the range of environments in the area. It is important to survey localised habitats within the FPP area. Communities with a high priority for conservation often occupy distinctive habitats (e.g. rocky knolls, poorly drained flats) or have fairly distinctive features (e.g. the white trunks of *E. viminalis* in *E. viminalis* wet forests contrast with the fibrous trunks of *E. obliqua* and *E. delegatensis*, the more widespread wet eucalypt forest dominants).

FPOs should try to identify a community occupying a small area (<1 ha) if:

- the forest in the small area is significantly different to the adjacent forest; or
- the forest community in the small area may be a priority community.

FPOs can subsume a community occupying a small area into the adjacent community if:

- the forest in the small area has obvious affinities to the forest community in the adjacent area (e.g. the same canopy dominants); and
- the forest community in the small area is not a priority community.

For RFA communities that only require additional conservation of oldgrowth occurrences, FPA should be notified for all oldgrowth patches exceeding 3 ha (including areas that extend beyond the FPP area).

There are particular problems in dealing with transitional vegetation and the RFA damp sclerophyll forest community (see discussion below). FPOs should take care not to confuse vegetation in transition zones with distinct communities.

When small areas of priority forest communities are referred to FPA, the advice given will be determined on a case-by-case basis. Factors that may be relevant include: requirements under the RFA and other policies; location within a coupe (e.g. whether adjacent to streamside reserve or in the middle of a proposed plantation); proposed silvicultural practices; presence of other values; and the local context of the community.

## Qualifications in the tables

There are many grey areas in classifying vegetation and determining conservation priorities. The tables give qualifications for some communities, when the conservation priority of the community will depend on particular circumstances. For example, some floristic communities can be allocated to more than one RFA community, depending on site characteristics (e.g. rock type) or co-occurring species. Grassy forests dominated by *E. globulus* are allocated to the RFA community “grassy *E. globulus* forest” (high conservation priority) if *E. pulchella* is sparse or absent, but are allocated to “*E. pulchella* - *E. globulus* - *E. viminalis* grassy shrubby forest” (low conservation priority) if *E. pulchella* is a subdominant species.

## More on community names and relationships

The systems of classifying floristic communities differ between forest types. This is because the classifications were undertaken by different researchers at different times. Most communities have an abbreviated name (used in the keys) and a more detailed name (used in the tables) that indicates some typical species or characteristics of the community. However, some stands of a particular community may not contain all the “typical” species given in the more detailed name of the community.

Most floristic communities can be readily allocated to RFA communities, but this is not always the case. Most RFA communities contain two or more floristic communities. In some cases, the RFA community names may seem inappropriate to describe some forest communities. For example, areas of wet sclerophyll forest dominated by *E. globulus* are included in the RFA community “*E. regnans* forest”.

## Dominance in forest communities

Accurate determination of the dominant canopy (overstorey) species and understorey characteristics is needed to classify communities. Most areas of forest contain one or more shrub layers below the canopy, and a ground layer of grasses, sedges, ferns or some combination of these. The dominant component of a vegetation layer is the species (or group of species) that supply most of the cover.

### Overstorey dominance

Identifying the dominant overstorey species is one of the first steps in keying out most forest communities. This can be difficult in forests containing more than one species of eucalypt. However, in most situations, one species is clearly dominant while the others are subdominant or minor. An example: *E. ovata* provides about 60% cover on a poorly drained flat with an understorey dominated by sedges; the flat also carries *E. amygdalina* and *E. viminalis*. The floristic community is sedgy *E. ovata* dry sclerophyll forest and the correlated RFA community is shrubby *E. ovata* - *E. viminalis* forest.

Two species occasionally occur as codominants, having about equal cover in the community. The community should be keyed out using both dominants as options. Botanical advice may be needed if one of those options is a priority community. An example: if *E. obliqua* and *E. viminalis* are codominant in a wet sclerophyll forest, the community can be identified as an *E. obliqua* wet sclerophyll forest community or an *E. viminalis* wet sclerophyll forest community. The latter community has a high priority for protection in all regions of Tasmania, and the operation needs to be referred to FPA. If neither community is a priority community (e.g. *E. obliqua* and *E. delegatensis* co-dominant in wet sclerophyll forest), the operation does not need to be referred to FPA unless other flora values are present. FPOs should exercise their own judgement (e.g. by taking account of associated vegetation and site characteristics) when allocating such forest to floristic and RFA communities.

### Understorey dominance

Within a broad forest type, some communities key out simply on the basis of their overstorey dominants. However, most floristic communities are keyed out by the presence or absence of understorey species (e.g. most wet sclerophyll forest communities) or by characteristics of the dominant understorey layer (e.g. most dry sclerophyll forest communities). For example, shrubs exceeding 2 m in height will be the most conspicuous understorey layer in a shrubby dry sclerophyll forest community. Grasses or sags are the most conspicuous understorey components in a grassy dry sclerophyll forest.

FPOs may need to make allowance if land uses or events have temporarily changed the nature of the understorey. For example, a recent fire may remove the shrub layer from a heathy forest, but if the vegetation in nearby areas or other evidence suggests that short shrubs are typically present, the community should be allocated to a heathy dry sclerophyll forest community. Section 2.6 gives more information on identifying the dominant understorey characteristics in dry sclerophyll communities.

### **Distinguishing eucalypt species**

Correct identification of eucalypt species is essential as they are the main tree species used to identify most dry sclerophyll, wet sclerophyll and mixed forest communities. Identification can sometimes be difficult because eucalypts hybridise readily. Seek botanical advice if you find unusual or outlying occurrences of eucalypts, as these may be genetically important.

The FPA website contains scanned images of Tasmanian eucalypts; a key to species and notes on distinguishing between some closely-related species (*E. viminalis* and *E. dalrympleana*; and *E. brookeriana* and *E. ovata*) associated with communities with a high priority for conservation. Useful references for identifying eucalypts are also listed in *Flora Technical Note 2*.

FPOs may need to collect material or take notes to determine the identity of a species. Bark characteristics, fruit, buds and adult and juvenile leaves can all be important for diagnosis. Juvenile leaves may be needed to identify some species (e.g. *E. viminalis* and *E. dalrympleana*).

### **Oldgrowth**

Oldgrowth forests have over-mature to senescent trees contributing over 30% of the crown cover to the overstorey, and have not been significantly affected by man-made disturbance. Fire does not preclude classification as oldgrowth, providing other oldgrowth characteristics are present. Oldgrowth forests generally contain a greater range of habitats than regrowth forests and consequently support a different (and generally more diverse) suite of species. Oldgrowth forest is discussed in *Flora Technical Note 7*.

Generally, oldgrowth forests have a higher conservation value than non-oldgrowth forests of the same community. Areas of oldgrowth forest, or areas containing oldgrowth trees, should be preferentially located in retained areas, if this is an option under the proposed silvicultural regime. Some RFA communities require additional protection for the oldgrowth component of the community only. For these communities, the practical minimum patch size that requires notification to FPA is 3 hectares (including areas extending beyond the coupe boundary).

## Transition zones

Transition zones often occur between adjacent forest types or adjacent forest communities, with vegetation of these zones being intermediate in structure and composition. Transition zones should be avoided when communities are being identified. Some forest communities (e.g. damp sclerophyll forest communities) are inherently intermediate in character and occupy relatively large areas - see below.

## Damp sclerophyll forest communities

Some eucalypt-dominated forests have an understorey with a similar proportion of wet sclerophyll species (e.g. broad-leaved shrubs and wet ferns) and dry sclerophyll species (e.g. narrow-leaved shrubs and grasses). An example of a damp sclerophyll understorey could include dogwood, blanket bush, prickly mo, prickly beauty, guitar plant and sagg. Such vegetation is sometimes described as damp sclerophyll forest. In this section, it should be keyed to its floristic community using the dry sclerophyll forest key (where it will generally key out as a shrubby dry sclerophyll community).

One of the RFA communities is *E. viminalis* - *E. ovata* - *E. amygdalina* - *E. obliqua* damp sclerophyll forest (DSC). The community has a damp sclerophyll understorey and *E. amygdalina* and/or *E. obliqua* are both prominent in the overstorey. *E. viminalis* and *E. ovata* may be present as subdominant or minor species or may dominate very small patches within a mosaic of forest dominated by *E. amygdalina* or *E. obliqua*. This community is mapped inconsistently on the RFA Forest Communities Map. On most sites mapped as DSC, the vegetation can be better allocated to other RFA communities (e.g. dry *E. obliqua* forest, tall *E. obliqua* forest, *E. amygdalina* forest on dolerite, shrubby *E. ovata* - *E. viminalis* forest).

## Inland *E. amygdalina* forest

The RFA community inland *E. amygdalina* forest (AI) was divided in 2005 into two distinct facies. Inland *E. amygdalina* - *E. viminalis* - *E. pauciflora* forest/woodland on Cainozoic deposits (AIC) is associated with Recent and Tertiary sediments (including ironstone lags) – it occurs mainly on private land and is listed as a Vulnerable community. It is abbreviated to Inland *E. amygdalina* forest on Cainozoic deposits in tables in Section 2.6. *E. amygdalina* forest on mudstone (AM) occurs locally on dry sites on Permian mudstone and Mathinna series substrate in Freycinet Region – it is not identified as a threatened community, though oldgrowth stands require protection on public land.

## Forest communities that are susceptible to *Phytophthora cinnamomi*

Some communities are very susceptible to the root rot pathogen *Phytophthora cinnamomi* because they:

- contain many species of susceptible plants, including threatened species;
- occur in warm, moist environments that are conducive to establishment of *Phytophthora*;
- occur in locations where spores can be transferred into uninfected sites by land use.

Forest communities that are highly susceptible to *Phytophthora* are identified (#) in the tables indicating conservation priorities of the different forest types (see column dealing with conservation status of the floristic community). Most are lowland dry sclerophyll forest communities – many are also Priority A communities. Several non-forest communities are also susceptible to *Phytophthora* – these should be referred to FPA if they are in operational areas (see below). Information on *Phytophthora* and its management in Tasmanian forests is given in Section 6 of this module and in *Flora Technical Note 8*.

## Non-forest vegetation

Native non-forest vegetation (e.g. moorland, heath, wetland and native grassland) may be associated with native forests (and sometimes plantations). Some of these vegetation types have a high priority for conservation, contain threatened species or are very susceptible to disturbance or disease. There are specific guidelines in the *Forest Practices Code* to avoid disturbance to localised environments (e.g. swamps, rocky knolls, streambanks) that often contain these vegetation types. The key on the following page will allow FPOs to identify broad non-forest vegetation types. Seek botanical advice in all cases where native non-forest vegetation will be affected by forestry operations.

## When to seek advice

This section of the module, and the FPP *Flora Evaluation Sheet*, indicates when botanical advice is needed because of the presence of particular communities in areas proposed for forestry operations. However, there is no shortage of grey areas in the natural world. Specialist advice should be sought if FPOs are uncertain about identification of communities or their conservation priority.

## KEY TO VEGETATION TYPES AND FOREST COMMUNITIES

Use the key to forest and non-forest types to identify the vegetation types present in the coupe, then go to the indicated section (forest types only) to identify the floristic communities. The table following the key will allow the floristic communities to be related to the RFA communities.

Each key should be followed through sequentially. A true/false decision should be made for each statement bearing the same number (e.g. ❶). If true, proceed to the next numbered statement immediately below (❷). If false (or there is some degree of doubt), proceed to the next statement of the same number (❶) in the key.

The keys are based on species or understorey types that will be familiar to most field workers. Understoreys are defined by their dominant species, although species typical of other vegetation types may be present. Information on species and other characteristics used to distinguish communities is provided in Module 1, *Flora Technical Note 2* and on the FPA Website.

Transitional vegetation may not key out easily. If the forest is intermediate between two recognisable floristic communities, assess the conservation priorities for both communities. Contact the FPA Botanist or Ecologist if a vegetation type or forest community does not key out.

### KEY TO FOREST TYPES


*Use when vegetation is dominated by trees exceeding 5 metres, or with potential to exceed 5 metres*

- ❶ **Eucalypts absent or less than 5% cover**
  - ❷ Myrtle, sassafras or celery-top pine dominant ..... Rainforest (go to 2.1)
  - ❷ Blackwood, tea-trees or paperbarks dominant ..... Swamp forest and related forest or scrub (go to 2.2)
  - ❷ Forest or tall scrub (5 to 8 m) with other species dominant ..... Other forest or scrub (go to 2.3)
- ❶ **Eucalypts present with greater than 5% cover**
  - ❷ Myrtle, sassafras or celery-top pine prominent as secondary trees or shrubs ..... Mixed forest (go to 2.4)
  - ❷ Understorey dominated by tall tea-trees or paperbarks ..... Wet sclerophyll forest (go to 2.5)
  - ❷ Understorey dominated by broad-leaved (soft-leaved) shrubs ..... Wet sclerophyll forest (go to 2.5)
  - ❷ Understorey dominated by an equal mixture of broad-leaved and narrow-leaved shrubs ..... Dry sclerophyll forest/woodland (go to 2.6)
  - ❷ Understorey dominated by grasses, sedges, heaths or narrow-leaved (hard-leaved) shrubs (often under 2 m in height) ..... Dry sclerophyll forest/woodland (go to 2.6)

### KEY TO NON-FOREST TYPES

*Seek advice from FPA in all cases if operations may affect native non-forest vegetation.*

- ❶ Dominated by shrubs over 2 m ..... Scrub
- ❶ Dominated by shrubs under 2 m, usually infertile or exposed sites ..... Heathland
- ❶ Dominated by sedges or buttongrass; low to high altitudes, often on the boundary of sedgy woodland or tea-tree scrub forest ..... Moorland/sedgeland
- ❶ Dominated by native grasses and sags; often herb-rich; generally fertile sites ..... Native grassland
- ❶ Dominated by *Sphagnum* moss; shrubs (e.g. tea-tree or richea) may be sparse or locally dense; often in high altitude soaks or drainage lines ..... *Sphagnum* peatland
- ❶ Aquatic vegetation or vegetation submerged seasonally, generally dominated by graminoids, herbs or succulent species ..... Wetland



Diagrams showing relationships between forest types and typical Tasmanian forest environments:

*A: Moderate rainfall site: soils of moderate fertility (e.g. dolerite); site varying in landform and fire history*

1 – shrubby *E. amygdalina* dry sclerophyll forest (exposed slope); 2 – grassy *E. amygdalina* dsf (exposed ridgeline); 3 – *E. delegatensis* wet sclerophyll forest (shaded slopes at higher altitudes); 4 – *E. delegatensis* mixed forest (humid slope, infrequently burnt); 5 – *E. obliqua* wsf and mixed forest (shaded slopes at lower altitudes); 6 – callidendrous rainforest (humid fire-shadow gully); 7 – *E. regnans* wsf (regrowth on humid site after wildfire or intensive logging).

*B: High rainfall site, low altitude: site varying greatly in soil fertility and drainage*

1 – heathy *E. amygdalina* dry sclerophyll forest (infertile substrate); 2 – *E. obliqua* wet sclerophyll forest (shaded slope); 3 – *E. obliqua* mixed forest (humid slope, infrequently burnt); 4 – callidendrous rainforest (humid, well-drained lower slope; fire-shadow site); 5 – *Leptospermum lanigerum* swamp forest (poorly-drained flat).

## 2.1 RAINFOREST COMMUNITIES

*Major References:* Jarman, Brown and Kantvilas (1984); Jarman, Kantvilas and Brown (1991); Neyland (1991); *Flora Technical Note 4*.

Tasmanian cool temperate rainforest is defined as vegetation with trees taller than 8 m, dominated by the following species: myrtle, deciduous beech, sassafras, leatherwood, horizontal, celery-top pine, King Billy pine, Huon pine, Cheshunt pine or pencil pine. Sassafras is the main dominant of rainforest patches in Freycinet Region, with myrtle and celery-top pine occurring locally on some sites. All rainforest in Freycinet Region occurs as relict stands, restricted to humid, fire-shadow sites. These mainly comprise moist, shaded gullies and creeklines, but small patches of rainforest are found on some coastal peaks and ridges where the vegetation strips moisture from sea-mist. Rainforest grades into mixed forest, wet sclerophyll forest and occasionally swamp forest, though transitions can be sharp across fire boundaries. All patches of rainforest in Freycinet Region have a high priority for conservation. Almost all patches have a callidendrous structure, but a few sites on Tasman and Forestier Peninsulas contain species typical of thamnian rainforest, and these sites are separately covered in the key.

### KEY TO RAINFOREST COMMUNITIES

#### ❶ Dominated by myrtle

- ❷ Ground ferns prominent
  - ❸ Musk common..... RAIN-C3.1
  - ❸ Musk sparse or absent
    - ❹ Celery-top pine or native laurel present..... RAIN-CT
    - ❹ Celery-top pine and native laurel absent..... RAIN-C1.1
- ❷ Ground ferns rare; woolly tea-tree or native pepper often present..... RAIN-C2.1

#### ❶ Dominated by sassafras

- ❷ Musk common ..... RAIN-C3.2
- ❷ Musk sparse or absent
  - ❹ Celery-top pine or native laurel present..... RAIN-CT
  - ❹ Celery-top pine and native laurel absent..... RAIN-C1.2

### RELICT RAINFOREST

Relict rainforest comprises isolated patches of rainforest that occur locally in humid or fire-shadow environments, outside the normal range of Tasmanian rainforest. All areas of rainforest in Freycinet Region are considered as relict rainforest. Eighty patches have been identified from the region, ranging in size from less than 1 ha to over 100 ha (e.g. Macgregor Peak on Forestier Peninsula and Big Sassy Creek in the Eastern Tiers). Grid references and further details of these sites are given in *Relict Rainforest in Eastern Tasmania* (Neyland 1991) and *Flora Technical Note 4*. Other patches of relict rainforest may occur in the region, most likely on sites with a PI type containing S or T.

Areas of relict rainforest that could be affected by forestry operations need to be referred to FPA. Typical prescriptions for protection of relict rainforest are given in *Flora Technical Note 4*.

**CONSERVATION PRIORITIES AND ATTRIBUTES OF RAINFOREST COMMUNITIES**

Note: \* – Community identified as Rare, Vulnerable or Endangered at a Statewide level through RFA processes

Floristic community code and name		RFA community code and name		Conservation priority		Distribution in Freycinet Region
				Floristic	RFA	
RAIN-C1.1 Callidendrous	<i>Nothofagus cunninghamii</i> - <i>Atherosperma moschatum</i> over <i>Dicksonia antarctica</i> and/or <i>Polystichum proliferum</i>	M+	Callidendrous & thamnic rainforest on fertile sites	A	N	Very moist, fire-shadow sites on S and E-facing slopes and gullies; tends to occur in higher rainfall areas than other rainforest communities in Freycinet Region (e.g. Tatnells Hill area, Tasman Peninsula).
RAIN-C1.2 Callidendrous	<i>Atherosperma moschatum</i> over <i>Dicksonia antarctica</i> - <i>Polystichum proliferum</i> - <i>Blechnum wattsii</i>	M+	Callidendrous & thamnic rainforest on fertile sites	A	N	Very moist, fire-shadow sites on S and E-facing slopes and gullies; most commonly associated with the upper reaches of deeply incised streams; often fringed by blackwood.
RAIN-C2.1 Callidendrous	<i>Nothofagus cunninghamii</i> - ( <i>Leptospermum lanigerum</i> ) over clear understorey or <i>Telopea truncata</i> and/or <i>Tasmannia lanceolata</i>	M+	Callidendrous & thamnic rainforest on fertile sites	A	N	Not recorded from Freycinet Region, but may occur at higher altitudes in localised areas of impeded drainage.
RAIN-C3.1 Callidendrous	<i>Nothofagus cunninghamii</i> - <i>Atherosperma moschatum</i> over <i>Olearia argophylla</i> with <i>Dicksonia antarctica</i> and/or <i>Polystichum proliferum</i>	M+	Callidendrous & thamnic rainforest on fertile sites	A	N	Moist fire-shadow sites and cloud-trapping upper slopes, generally with SE aspect, on Tasman and Forestier Peninsulas.
RAIN-C3.2 Callidendrous	<i>Atherosperma moschatum</i> over <i>Olearia argophylla</i> with <i>Dicksonia antarctica</i> and/or <i>Polystichum proliferum</i>	M+	Callidendrous & thamnic rainforest on fertile sites	A	N	The most widespread of the Freycinet Region rainforest communities. Mainly found on fire-shadow sites within wet eucalypt forests of the Eastern Tiers (e.g. Big Sassy Creek), Wielangta (Sandspit River, Mt Walter), and Tasman and Forestier Peninsulas (e.g. Macgregor Peak).
RAIN-CT Callidendrous – thamnic intermediate	varies	M+	Callidendrous & thamnic rainforest on fertile sites	A	N	Very localised on moist fire-shadow sites and cloud-trapping upper slopes, on Tasman and Forestier Peninsulas (e.g. Tatnells Hill, Balts Spur Road).

## 2.2 SWAMP FOREST AND RELATED FOREST OR SCRUB COMMUNITIES

*Major Reference:* Pannell (1992)

Swamp forests have a closed canopy of blackwood, tea-trees or paperbarks, and typically occupy flat, poorly drained sites. Riparian blackwood forest contains some typical wet sclerophyll species (e.g. dogwood, cheesewood, stinkwood and bracken) that are not found in swamp forests. They occur on better drained sites adjacent to rivers and creeks, and extend to shaded slopes.

Swamp forests and related forest types have a very restricted distribution in Freycinet Region. They occur locally on moderate to high rainfall sites with impeded drainage, mainly in the Eastern Tiers, Wielangta area and on Forestier and Tasman Peninsulas. All sites are potentially important for conservation.

### KEY TO SWAMP FOREST AND RELATED FOREST OR SCRUB COMMUNITIES

- ❶ **Blackwood, paperbark or woolly tea-tree dominant; understorey open with ferns or cutting grass common; mainly found on poorly drained sites**
  - ❷ Swamp paperbark common; near-coastal sites .....SWAMP-C2
  - ❷ Swamp paperbark absent
    - ❸ Sassafras common..... SWAMP-A2
    - ❸ Sassafras sparse or absent
      - ❹ Woolly tea-tree common; scented paperbark present; cutting sedge absent..... SWAMP-A1
      - ❹ Woolly tea-tree sparse or absent..... SWAMP-A3
- ❶ **Blackwood dominant or codominant; dogwood, cheesewood, musk, cathead fern or bracken prominent in understorey; mainly found on better drained flats, riparian sites and slopes**
  - ❷ Silver wattle common; woolly tea-tree often present; dogwood sparse or absent ..... SWAMP-D2
  - ❷ Silver wattle and woolly tea-tree sparse or absent; dogwood common ..... SWAMP-D4
- ❶ **Woolly tea-tree dominant and montane sites (usually above 500 m)**
  - ❷ Myrtle sparse or absent
    - ❸ Silver wattle codominant or common .....SWAMP-E1
    - ❸ Silver wattle sparse or absent .....SWAMP-E1
  - ❷ Myrtle codominant or common (silver wattle sparse or absent).....SWAMP-E2
- ❶ **Tea-tree (manuka) usually dominant or codominant; usually lowland sites..... SWAMP-F1**

**CONSERVATION PRIORITIES AND ATTRIBUTES OF SWAMP FOREST AND RELATED FOREST OR SCRUB COMMUNITIES**

Note: \* – Community identified as Rare, Vulnerable or Endangered at a Statewide level through RFA processes

Floristic community code and name		Qualification	RFA community code and name		Conservation priority		Distribution in Freycinet Region
					Floristic	RFA	
SWAMP-A1	Depauperate callidendrous swamp forest		BF	<i>Acacia melanoxylon</i> forest on flats	A	N	May occur locally on poorly drained flats (e.g. Tasman Peninsula).
SWAMP-A2	Callidendrous sassafras swamp forest		BF	<i>Acacia melanoxylon</i> forest on flats	A	N	May occur locally on poorly drained flats (e.g. Tasman Peninsula).
SWAMP-A3	Depauperate callidendrous fern swamp forest		BF	<i>Acacia melanoxylon</i> forest on flats	A	N	May occur locally on poorly drained flats, less recently disturbed than SWAMP-A1 or A2.
SWAMP-C2	Depauperate coastal paperbark swamp forest		ME	<i>Melaleuca ericifolia</i> coastal swamp forest	A	Y*	Local on poorly drained coastal sites in northern part of region (e.g. Templestowe Lagoon).
SWAMP-D2	Riparian blackwood/ wattle forest	On flats	BF	<i>Acacia melanoxylon</i> forest on flats	A	N	Not recorded from region, but may occur locally in gullies and on moist fertile lowland flats, where drainage is not too impeded.
		On rises	BR	<i>Acacia melanoxylon</i> forest on rises			
SWAMP-D4	Riparian blackwood/ dogwood forest	On flats	BF	<i>Acacia melanoxylon</i> forest on flats	A	N	Local in gullies and on moist fertile lowland flats, where drainage is not too impeded.
		On rises	BR	<i>Acacia melanoxylon</i> forest on rises			
SWAMP-E1	Depauperate montane tea-tree forest	Silver wattle common	SI	<i>Acacia dealbata</i> forest	B	N	Local on flats, often disturbed; mainly in dolerite uplands.
		Silver wattle sparse or absent	L	<i>Leptospermum</i> species - <i>Melaleuca squarrosa</i> swamp forest	B	N	Local on flats with impeded drainage, mainly in dolerite uplands (e.g. Mt Foster).
SWAMP-E2	Montane myrtle tea-tree forest		M+	Callidendrous & thamnic rainforest on fertile sites	A	N	Not recorded from region, but may occur on poorly drained upland sites that have not been burnt or disturbed for a long period.
SWAMP-F1	Depauperate coastal paperbark swamp forest		L	<i>Leptospermum</i> species - <i>Melaleuca squarrosa</i> swamp forest	A	N	Local on poorly drained sites, often in coastal areas on infertile substrate (e.g. Mt Fortescue area), but may also occurs locally on dolerite in the Eastern Tiers.

## 2.3 OTHER FOREST OR SCRUB COMMUNITIES

*Major Reference:* Kirkpatrick, Barker, Brown, Harris and Mackie (1995)

This section covers an array of forest communities that are not dominated by rainforest species, eucalypts, blackwood, tea-trees or paperbarks. They occupy a wide environmental range, from humid sites capable of supporting rainforest, to dry rocky gorges. Most of these communities occur as localised patches in other forest types. Examples include small stands (or groves) of native olive associated with rocky sites in wet sclerophyll forest; and she-oak forests on very dry hillsides.

Most of the communities have a high priority for conservation, because of their localised distribution. Some are associated with threatened species. The exception is silver wattle (*Acacia dealbata*) forest, which is found locally where inadequate regeneration of eucalypts has followed land clearing, wildfire or logging.

### KEY TO OTHER FOREST OR SCRUB COMMUNITIES

*Note:* These communities may have a sparse (<5%) cover of eucalypts or other tree species.

- ❶ Bull-oak dominant ..... DRY-LIT
- ❶ She-oak dominant
  - ❷ Non-coastal environments ..... DRY-VERT-inland
  - ❷ Coastal environments ..... DRY-VERT-coastal
- ❶ Dogwood (native pear), pinkwood or currajong dominant ..... OTHER-03
- ❶ Oyster Bay pine dominant ..... OTHER-05
- ❶ Blanket bush or musk dominant ..... OTHER-06
- ❶ Native olive (dorrel) dominant ..... OTHER-07
- ❶ Silver wattle dominant ..... OTHER-10
- ❶ Yellow bottlebrush (*Callistemon pallidus*) dominant ..... OTHER-11
- ❶ Black wattle dominant, often with prickly box and scattered *E. viminalis* ..... OTHER-12

## CONSERVATION PRIORITIES AND ATTRIBUTES OF OTHER FOREST OR SCRUB COMMUNITIES

Note: \* – Community identified as Rare, Vulnerable or Endangered at a Statewide level through RFA processes

Floristic community code and name		Qualification	RFA community code and name		Cons. priority		Distribution in Freycinet Region
					Floristic	RFA	
DRY-LIT	<i>Allocasuarina littoralis</i> woodland		check	check	A	check	Local on dry siliceous sites on sediments and granite, mainly in the north of Freycinet Region.
DRY-VERT coastal	Coastal <i>Allocasuarina verticillata</i> low forest		AV	<i>Allocasuarina verticillata</i> forest	A	Y	Rocky coastal sites, on a range of substrates (e.g. Lime Bay, Cape Bernier, Freycinet Peninsula).
DRY-VERT inland	Inland <i>Allocasuarina verticillata</i> low forest		AV	<i>Allocasuarina verticillata</i> forest	A	Y	Local on drought-prone dolerite sites (e.g. Cherry Tree Hill) - may displace grassy eucalypt woodlands in some areas.
OTHER-03	<i>Pomaderris apetala</i> - <i>Beyeria viscosa</i> - <i>Asterotrichion discolor</i> closed forest/scrub	Site disturbed by heavy logging or clearing	check	check	check	check	Occasional where very poor eucalypt regeneration has occurred following logging or clearing of wet forest.
		Not as above	NP	<i>Notelaea ligustrina</i> and/or <i>Pomaderris apetala</i> closed forest	A	Y*	Shaded creeks and gullies in drier areas, but also localised patches on moist slopes (e.g. Wielangta area, Kafoozalum Ridge).
OTHER-06	<i>Bedfordia salicina</i> - <i>Olearia argophylla</i> closed forest/ scrub	Site disturbed by heavy logging or clearing	check	check	check	check	Occasional where very poor eucalypt regeneration has occurred following logging or clearing of wet forest.
		Not as above	—	None appropriate	A	—	Shaded creeks and gullies in drier areas, but also localised on moist slopes. Mainly on dolerite, from uplands (e.g. Mt Tooms area) to steep coastal slopes (e.g. Cape Bernier).
OTHER-05	<i>Callitris rhomboidea</i> - <i>Bedfordia salicina</i> closed forest/ scrub		CR	<i>Callitris rhomboidea</i> forest	A	Y*	Locally common on rocky dolerite sites, protected from frequent fire, including steep coastal slopes (e.g. Tasman Peninsula), ridges and gullies (e.g. Castle Sugarloaf).
OTHER-07	<i>Notelaea ligustrina</i> closed forest		NP	<i>Notelaea ligustrina</i> and/or <i>Pomaderris apetala</i> closed forest	A	Y*	Occurs locally on rocky dolerite sites, generally protected from frequent fire, including steep creeks (e.g. Pigeon Creek) and benches (e.g. Douglas-Apsley NP).
OTHER-10	<i>Acacia dealbata</i> forest		SI	<i>Acacia dealbata</i> forest	B	N	Local on sites reverting to forest (with poor eucalypt regeneration) after clearing or successive fires (e.g. Nugent).
OTHER-11	<i>Callistemon pallidus</i> closed forest		—	None appropriate	A	—	Local on dolerite ridges and upper slopes that have not been burnt for a long time (e.g. Little Swanport Hill).
OTHER-12	<i>Acacia mearnsii</i> forest/woodland		—	None appropriate	A	—	Typically occurs as grassy woodland on dry dolerite hills and slopes (e.g. Swansea, Cranbrook, Sorell). Sometimes occurs as a scrub community invading paddocks.

## 2.4 MIXED FOREST COMMUNITIES

*Major Reference:* Kirkpatrick, Peacock, Cullen and Neyland (1988)

Mixed forest comprises vegetation with an understorey of rainforest species and an overstorey of eucalypts that becomes sparse as the forest approaches maturity. Often only one species of eucalypt is present, with trees frequently exceeding 50 m in mature forest. The eucalypts tend to be even-aged, and are usually of one or two age classes, which relate to period since fire or other major disturbance. Mixed forests have a minimum eucalypt canopy cover of 5% - if eucalypt cover is less than 5% the forest is considered as rainforest. Mixed forests represent a transition (in space or time) between the rainforests and the wet sclerophyll forests into which they grade.

Mixed forests in Freycinet Region are largely restricted to long-unburnt sites on moist, shaded slopes and gullies. They occur locally in the Eastern Tiers, Wielangta area and Forestier and Tasman Peninsulas. Most communities occur in formal reserves, but some communities are very localised and of considerable scientific interest.

### KEY TO MIXED FOREST COMMUNITIES

- ❶ Dominated by *E. delegatensis*
  - ❷ Celery-top pine, goldeywood, native plum, dragon heath or native laurel common; musk or manfern sparse or absent..... WET-DEL1011
  - ❷ Celery-top pine, goldeywood, native plum, dragon heath and native laurel sparse or absent; musk or manfern often common
    - ❸ Cutting grass common
      - ❹ Stinkwood, goldeywood or bracken common ..... WET-DEL0111
      - ❹ Stinkwood, goldeywood and bracken sparse or absent ..... WET-DEL1001
    - ❸ Cutting grass sparse or absent
      - ❹ Stinkwood or goldeywood common..... WET-DEL0111
      - ❹ Stinkwood and goldeywood sparse or absent
        - ❺ Myrtle present ..... WET-DEL1000
        - ❺ Myrtle absent ..... WET-DEL0110
- ❶ Dominated by *E. johnstonii*..... WET-JOHN1
- ❶ Dominated by *E. obliqua*
  - ❷ Cutting grass, prickly mo or goldeywood common; epiphytic ferns sparse or absent..... WET-OB101
  - ❷ Cutting grass, prickly mo and goldeywood sparse or absent; epiphytic ferns often common ..... WET-OB1000
- ❶ Dominated by *E. regnans*
  - ❷ Silver wattle or blackwood present; dogwood common; epiphytic ferns sparse or absent..... WET-REG101
  - ❷ Silver wattle and blackwood absent; dogwood sparse or absent; epiphytic ferns common ..... WET-REG110
- ❶ Dominated by *E. urnigera*..... WET-URN1
- ❶ Dominated by *E. viminalis* ..... WET-VIM111

CONSERVATION PRIORITIES AND ATTRIBUTES OF MIXED FOREST COMMUNITIES

Note: \* – Community identified as Rare, Vulnerable or Endangered at a Statewide level through RFA processes

Floristic community code and name		Qualification	RFA community code and name		Cons. priority		Distribution in Freycinet Region
					Floristic	RFA	
WET-DAL10	<i>E. dalrympleana</i> - <i>Tasmannia lanceolata</i> - <i>Dicksonia antarctica</i> mixed forest	Make sure dominant not <i>E. viminalis</i> – Contact FPA if unsure	DT	Tall <i>E. delegatensis</i> forest	A	N	May occur locally on shaded sites in upland areas of the Eastern Tiers.
WET-DEL0110	<i>E. delegatensis</i> - <i>Atherosperma moschatum</i> - <i>Olearia argophylla</i> wsf/mixed forest		DT	Tall <i>E. delegatensis</i> forest	np	N	Moist, shaded slopes, gullies and drainage headwaters on fertile sites in upland areas (e.g. Big Sassy Creek).
WET-DEL0111	<i>E. delegatensis</i> - <i>Zieria arborescens</i> - <i>Hydrocotyle sibthorpioides</i> wsf/mixed forest		DT	Tall <i>E. delegatensis</i> forest	B	N	May occur locally on moist upland sites of relatively low nutrient status (e.g. sandstone) on Tasman Peninsula.
WET-DEL1000	<i>E. delegatensis</i> - <i>Nothofagus cunninghamii</i> - <i>Grammitis billardierei</i> mixed forest		DT	Tall <i>E. delegatensis</i> forest	np	N	Moist, shaded slopes, drainage headwaters and gullies on more humid or fire-protected sites than DEL0110. Often associated with relict rainforest.
WET-DEL1011	<i>E. delegatensis</i> - <i>Monotoca glauca</i> - <i>Hymenophyllum rarum</i> mixed forest		DT	Tall <i>E. delegatensis</i> forest	B	N	Localised on less fertile and poorly drained sites in high rainfall parts of the region (e.g. Tatnells Hill area).
WET-JOHN1	<i>E. johnstonii</i> mixed forest		SU	<i>E. subcrenulata</i> forest	A	N	Very localised on fire-protected margins of poorly drained sites on Tasman Peninsula.
WET-OB1000	<i>E. obliqua</i> - <i>Nothofagus cunninghamii</i> - <i>Polystichum proliferum</i> - <i>Hymenophyllum flabellatum</i> mixed forest		OT	Tall <i>E. obliqua</i> forest	np	N	Locally common on fire-protected gullies and shaded slopes at lower altitudes (e.g. Meetus Falls, Mayson River). Often associated with relict rainforest.
WET-OB101	<i>E. obliqua</i> - <i>Nothofagus cunninghamii</i> - <i>Monotoca glauca</i> mixed forest		OT	Tall <i>E. obliqua</i> forest	np	N	Distribution as for WET-OB1000; may be an early successional phase of this community.
WET-REG101	<i>E. regnans</i> - <i>Atherosperma moschatum</i> - <i>Acacia dealbata</i> - <i>Olearia argophylla</i> wsf/mixed forest		R	<i>E. regnans</i> forest	np	N	Local in moist, fire-protected gullies and slopes (e.g. Sandspit River), but also occurs on upper slopes subject to sea-mist (e.g. Mt Walter). May be associated with relict rainforest.
WET-REG110	<i>E. regnans</i> - <i>Nothofagus cunninghamii</i> - <i>Atherosperma moschatum</i> mixed forest		R	<i>E. regnans</i> forest	A	N	Very local on more humid or fire-protected sites than REG101. Generally associated with relict rainforest.

**Forest Botany Manual: Module 4 – Freycinet Region**

Floristic community code and name		Qualification	RFA community code and name		Conservation priority		Distribution in Freycinet Region
					Floristic	RFA	
WET-URN1	<i>E. urnigera</i> wsf/mixed forest		C	<i>E. coccifera</i> forest	A	N	Recorded only from shaded upper slopes of Mt Maria
WET-VIM111	<i>E. viminalis</i> - <i>Nothofagus cunninghamii</i> - <i>Atherosperma moschatum</i> - <i>Dicksonia antarctica</i> mixed forest		VW	<i>E. viminalis</i> wet forest	A	Y*	Very local in humid, fire protected sites on fertile flats (e.g. Anglers Creek, Tooms Lake FR).

## 2.5 WET SCLEROPHYLL FOREST COMMUNITIES

*Major Reference:* Kirkpatrick, Peacock, Cullen and Neyland (1988)

Wet sclerophyll forests are typically dominated by eucalypts and have an understorey dominated by broad-leaved (soft-leaved) shrubs. Trees in mature forest generally exceed 40 m. Wet sclerophyll forests typically contain only one or two eucalypt age classes - these relate to period since fire or other major disturbance (including intensive logging and regeneration burning). Often only one species of eucalypt is present. The shrub layer is generally dense, preventing continuous regeneration of shade-intolerant species such as eucalypts. Ferns are often prominent in the ground layer.

Wet sclerophyll communities are relatively widespread in higher rainfall parts of Freycinet Region (e.g. Tasman Peninsula, Forestier Peninsula, Wielangta area) and in moist, cool upland environments (e.g. Eastern Tiers). In drier parts of the region, they are largely restricted to riverine corridors and moist, shaded slopes and gullies. Wet sclerophyll forest grades into mixed forest (as rainforest species become more prominent in the understorey) and into dry sclerophyll forest (often through a damp sclerophyll transition zone) as sites become more exposed to drought and fire. However, on some sites the transition between wet sclerophyll and dry sclerophyll forests is remarkably sharp. Most wet sclerophyll forest communities are well reserved, but some localised communities (e.g. wet sclerophyll forests dominated by *E. viminalis* or *E. brookeriana*) have a high priority for conservation.

### KEY TO WET SCLEROPHYLL FOREST COMMUNITIES

- ❶ Dominated by *E. amygdalina* ..... WET-AM1
- ❶ Dominated by *E. brookeriana*
  - ❷ Blanket bush, musk or bracken common; prickly mo and sword sedge sparse or absent ..... WET-BR11
  - ❷ Blanket bush, musk and bracken sparse or absent; prickly mo or sword sedge common ..... WET-BR2
- ❶ Dominated by *E. dalrympleana*
  - ❷ Guitar plant, prickly beauty or sagg common; manfern and musk absent ..... WET-DAL00
  - ❷ Guitar plant, prickly beauty and sagg sparse or absent; manfern or musk often present.... WET-DAL01
- ❶ Dominated by *E. delegatensis*
  - ❷ Guitar plant or prickly beauty common
    - ❸ Mountain currant, waratah, native pepper or tussock grass common
      - ❹ Dogwood, dwarf musk, viscid daisy bush or flax lily common..... WET-DEL0010
      - ❹ Dogwood, dwarf musk, viscid daisy bush and flax lily sparse or absent ..... WET-DEL0011
    - ❸ Mountain currant, waratah, native pepper and tussock grass sparse or absent
      - ❹ Dogwood, manfern or cutting grass common
        - ❺ Stinkwood or goldeywood common; blanket bush absent ..... WET-DEL0111
        - ❺ Stinkwood and goldeywood sparse or absent; blanket bush often present.... WET-DEL0100
      - ❹ Dogwood, manfern and cutting grass sparse or absent
        - ❺ Silver wattle or native currant common; blackwood, kangaroo fern, dwarf musk or dolly bush sparse or absent ..... WET-DEL0000
        - ❺ Silver wattle and native currant sparse or absent; blackwood, kangaroo fern, dwarf musk and dolly bush common ..... WET-DEL0001
  - ❷ Guitar plant and prickly beauty sparse or absent
    - ❸ Cutting grass or sword sedge common
      - ❹ Woolly tea-tree common; goldeywood or stinkwood sparse or absent..... WET-DEL3
      - ❹ Woolly tea-tree sparse or absent; goldeywood or stinkwood common..... WET-DEL0111
    - ❸ Cutting grass or sword sedge sparse or absent
      - ❹ Dwarf musk, viscid daisy bush or bracken common; drier sites..... WET-DEL0101
      - ❹ Dwarf musk, viscid daisy bush or bracken sparse or absent; moister sites ..... WET-DEL0110

- ❶ Dominated by *E. globulus*
  - ❷ Tussock grass (*Poa*), sagg or prickly mo common
    - ❸ Cutting grass or musk present; viscid daisy bush sparse or absent ..... WET-GLOB0101
    - ❸ Cutting grass and musk sparse or absent; viscid daisy bush often present..... WET-GLOB1
  - ❷ Tussock grass (*Poa*), sagg and prickly mo sparse or absent
    - ❸ Manfern, kangaroo fern or epiphytic ferns present; cutting grass absent ..... WET-GLOB001
    - ❸ Manfern, kangaroo fern and epiphytic ferns absent; cutting grass often present
      - ❹ Blanket bush or pinkwood common, tea-tree and paperbark sparse or absent ..... WET-GLOB0100
      - ❹ Blanket bush and pinkwood sparse or absent; tea-tree or paperbark often present ..... WET-GLOB0101
- ❶ Dominated by *E. johnstonii* ..... WET-JOHN2
- ❶ Dominated by *E. obliqua*
  - ❷ Guitar plant or prickly beauty common; drier sites ..... WET-OB010
  - ❷ Guitar plant and prickly beauty sparse or absent; moister sites
    - ❸ Paperbark, tea-tree, sword sedge or cutting grass common; musk, silver wattle, blanket bush and pinkwood sparse or absent ..... WET-OB0111
    - ❸ Paperbark, tea-tree, sword sedge and cutting grass sparse or absent, musk silver wattle, blanket bush or pinkwood common..... WET-OB0110
- ❶ Dominated by *E. ovata*
  - ❷ Understorey dominated by paperbarks or tea-trees; cutting grass or sword sedge usually common ..... WET-OV00
  - ❷ Understorey dominated by broad-leaved shrubs; cutting grass and sword sedge usually sparse or absent ..... WET-OV01
- ❶ Dominated by *E. regnans*
  - ❷ Bracken, cutting grass or sword sedge common; (myrtle or sassafras generally absent)
    - ❸ Dwarf musk, prickly mo, stinkwood or goldeywood common ..... WET-REG1000
    - ❸ Dwarf musk, prickly mo, stinkwood and goldeywood sparse or absent ..... WET-REG1001
  - ❷ Bracken, cutting grass and sword sedge sparse or absent; (myrtle or sassafras often present) ..... WET-REG101
- ❶ Dominated by *E. sieberi* ..... WET-SIEB01
- ❶ Dominated by *E. urnigera* ..... WET-URN2
- ❶ Dominated by *E. viminalis*
  - ❷ Paperbark or tea-tree common; *E. ovata* often present; poorly drained sites..... WET-VIM2
  - ❷ Paperbark or tea-tree sparse or absent; *E. ovata* absent; well drained sites
    - ❸ Native cherry, guitar plant or little prickly common; drier sites ..... WET-VIM0011
    - ❸ Native cherry, guitar plant and little prickly sparse or absent; moister sites
      - ❹ Prickly mo or dolly bush common; blanket bush and manfern sparse or absent. WET-VIM0100
      - ❹ Prickly mo and dolly bush sparse or absent; blanket bush or manfern common. WET-VIM0101

**CONSERVATION PRIORITIES AND ATTRIBUTES OF WET SCLEROPHYLL FOREST COMMUNITIES**

Note: \* – Community identified as Rare, Vulnerable or Endangered at a Statewide level through RFA processes

Floristic community code and name		Qualification	RFA community code and name		Conservation priority		Distribution in Freycinet Region
					Floristic	RFA	
WET-AM1	<i>E. amygdalina</i> - <i>E. viminalis</i> - <i>Lomandra longifolia</i> wsf		AD	<i>E. amygdalina</i> forest on dolerite	np	N	Moderately shaded slopes on dolerite in the Eastern Tiers; sites often rocky.
WET-BR11	<i>E. brookeriana</i> - <i>E. obliqua</i> - <i>Bedfordia salicina</i> wsf		BA	<i>E. brookeriana</i> wet forest	A	Y*	Locally common in well drained gully headwaters in the Eastern Tiers (e.g. Rocka Rivulet), and also occurs on flats in the SE of the region (e.g. Wielangta area).
WET-BR2	<i>E. brookeriana</i> - <i>Leptospermum</i> species - <i>Lepidosperma elatius</i> wsf		BA	<i>E. brookeriana</i> wet forest	A	Y*	Local on poorly drained flats in the Wielangta area, and possibly Forestier Peninsula and Tasman Peninsula.
WET-DAL00	<i>E. dalrympleana</i> / <i>E. delegatensis</i> - <i>Lomatia tinctoria</i> wsf	Make sure dominant is not <i>E. viminalis</i> – contact FPA if unsure	DT	Tall <i>E. delegatensis</i> forest	B	N	Occurs locally in shaded upland areas of the Eastern Tiers, in association with <i>E. delegatensis</i> forest.
WET-DAL01	<i>E. dalrympleana</i> - <i>Pomaderris apetala</i> - <i>Bedfordia salicina</i> wsf		DT	Tall <i>E. delegatensis</i> forest	B	N	
WET-DEL0000	<i>E. delegatensis</i> - <i>Bedfordia salicina</i> - <i>Lomatia tinctoria</i> wsf		DT	Tall <i>E. delegatensis</i> forest	np	N	Widespread on upland dolerite slopes and tier surfaces.
WET-DEL0001	<i>E. delegatensis</i> - <i>Acacia melanoxylon</i> - <i>Bedfordia salicina</i> wsf		DT	Tall <i>E. delegatensis</i> forest	np	N	Local on high altitude, rocky dolerite sites (e.g. Mt Foster).
WET-DEL0010	<i>E. delegatensis</i> - <i>Olearia phlogopappa</i> - <i>Olearia viscosa</i> wsf		DT	Tall <i>E. delegatensis</i> forest	B	N	Local on rocky high altitude slopes and boulder fields (e.g. Mt Maria).
WET-DEL0011	<i>E. delegatensis</i> - <i>Telopea truncata</i> wsf		DT	Tall <i>E. delegatensis</i> forest	np	N	Considered a relatively wet expression of shrubby <i>E. delegatensis</i> dry sclerophyll forest. Recorded from Mt Foster.
WET-DEL0100	<i>E. delegatensis</i> - <i>E. viminalis</i> - <i>Acacia melanoxylon</i> wsf		DT	Tall <i>E. delegatensis</i> forest	np	N	Common on shaded and sometimes disturbed sites at the lower altitudinal limit of <i>E. delegatensis</i> (about 300 m); often associated with <i>E. obliqua</i> wet sclerophyll forest.

**Forest Botany Manual: Module 4 – Freycinet Region**

Floristic community code and name		Qualification	RFA community code and name		Conservation priority		Distribution in Freycinet Region
					Floristic	RFA	
WET-DEL0101	<i>E. delegatensis</i> - <i>E. obliqua</i> - <i>Acaena novae-zelandiae</i> wsf		DT	Tall <i>E. delegatensis</i> forest	np	N	Occurs at the lower altitude limit of <i>E. delegatensis</i> , commonly forming a transition zone with <i>E. obliqua</i> wet sclerophyll forest.
WET-DEL0110	<i>E. delegatensis</i> - <i>Atherosperma moschatum</i> - <i>Olearia argophylla</i> wsf/mixed forest		DT	Tall <i>E. delegatensis</i> forest	np	N	Relatively common in upland drainage headwaters, creeklines and shaded slopes, often grading into mixed forest (e.g. Big Sassy Creek, and Meetus Falls area).
WET-DEL0111	<i>E. delegatensis</i> - <i>Zieria arborescens</i> - <i>Hydrocotyle sibthorpioides</i> wsf/mixed forest		DT	Tall <i>E. delegatensis</i> forest	B	N	Localised on moist upland sites of relatively low nutrient status (e.g. sandstone) on Tasman Peninsula.
WET-DEL3	<i>E. delegatensis</i> - <i>Leptospermum lanigerum</i> - <i>Gahnia grandis</i> wsf		DT	Tall <i>E. delegatensis</i> forest	B	N	Drainage headwaters and poorly drained saddles, in higher rainfall parts of the region (e.g. Tatnells Hill, Mt Fortescue).
WET-GLOB001	<i>E. globulus</i> - <i>Dicksonia antarctica</i> - <i>Ctenopteris heterophylla</i> wsf		R	<i>E. regnans</i> forest	B	N	Sheltered slopes and gullies, mainly on dolerite (e.g. Sandspit River). <i>E. obliqua</i> or <i>E. regnans</i> often also present.
WET-GLOB0100	<i>E. globulus</i> - <i>Bedfordia salicina</i> - <i>Beyeria viscosa</i> wsf		R	<i>E. regnans</i> forest	B	N	Common on shaded slopes and gully flanks (e.g. Blue Gum Spur, Wielangta, Bluemans Creek).
WET-GLOB0101	<i>E. globulus</i> - <i>Acacia dealbata</i> - <i>Acacia melanoxylon</i> - <i>Cassinia aculeata</i> wsf		R	<i>E. regnans</i> forest	B	N	Local on shaded slopes and gully flanks (e.g. Lower Marsh Creek), sometimes extending into poorly drained flats (e.g. Fortescue Road).
WET-GLOB1	<i>E. globulus</i> - <i>Poa labillardierei</i> - <i>Hypochoeris radicata</i> wsf		GG	Grassy <i>E. globulus</i> forest	A	Y*	Moister aspects of dry hills and on wetter sites which are grazed or burnt frequently. May form transition zone between wet and dry sclerophyll forest (e.g. Maria Island).
WET-JOHN2	<i>E. johnstonii</i> wsf		SU	<i>E. subcrenulata</i> forest	A	N	Local on moist benches on sandstone and dolerite; drainage often impeded; mainly on Tasman Peninsula (e.g. Mt Fortescue, Tatnells Hill).
WET-OB010	<i>E. obliqua</i> - <i>Olearia lirata</i> - <i>Pultenaea juniperina</i> wsf		OT	Tall <i>E. obliqua</i> forest	np	N	Common on fairly dry sites at lower altitudes (e.g. Wielangta area); often forms an intermediate community between wet sclerophyll and dry sclerophyll forests.

**Forest Botany Manual: Module 4 – Freycinet Region**

Floristic community code and name		Qualification	RFA community code and name		Conservation priority		Distribution in Freycinet Region
					Floristic	RFA	
WET-OB0110	<i>E. obliqua</i> - <i>Acacia dealbata</i> - <i>Olearia argophylla</i> wsf		OT	Tall <i>E. obliqua</i> forest	np	N	The most widespread wet sclerophyll forest community in the region; found on a wide range of substrates.
WET-OB0111	<i>E. obliqua</i> - <i>Melaleuca squarrosa</i> - <i>Monotoca glauca</i> wsf		OT	Tall <i>E. obliqua</i> forest	np	N	Wet or poorly drained sites; particularly along creeks and on river flats in wetter parts of the region (e.g. Forestier Peninsula, Tasman Peninsula).
WET-OV00	<i>E. ovata</i> - <i>Leptospermum lanigerum</i> wsf		OV	Shrubby <i>E. ovata</i> - <i>E. viminalis</i> forest	A	Y*	Local on poorly drained lowland flats that have escaped fire for a long period.
WET-OV01	<i>E. ovata</i> - <i>Acacia dealbata</i> - <i>Pomaderris apetala</i> wsf		OV	Shrubby <i>E. ovata</i> - <i>E. viminalis</i> forest	A	Y*	Local on shaded or fertile lowland flats where drainage less impeded than in most <i>E. ovata</i> -dominated communities.
WET-REG1000	<i>E. regnans</i> - <i>E. obliqua</i> - <i>Pomaderris apetala</i> - <i>Olearia lirata</i> wsf		R	<i>E. regnans</i> forest	B	N	Local on shaded slopes and gully flanks, in transition zone between <i>E. obliqua</i> wet sclerophyll forest and wetter <i>E. regnans</i> communities. Mainly occurs as a regrowth community.
WET-REG1001	<i>E. regnans</i> - <i>Acacia dealbata</i> - <i>Pomaderris apetala</i> wsf		R	<i>E. regnans</i> forest	B	N	Sheltered slopes and gullies in drier areas (e.g. Mt Morrison) but extending to drier sites in high rainfall areas. Mainly occurs as a regrowth community.
WET-REG101	<i>E. regnans</i> - <i>Atherosperma moschatum</i> - <i>Acacia dealbata</i> - <i>Olearia argophylla</i> wsf/mixed forest		R	<i>E. regnans</i> forest	np	N	Local on moist, shaded slopes and gullies in high rainfall areas (e.g. Tasman Peninsula, Wielangta) where it grades into mixed forest.
WET-SIEB01	<i>E. sieberi</i> - <i>Olearia argophylla</i> - <i>Coprosma quadrifida</i> wsf/mixed forest	Substrate granite	SG	<i>E. sieberi</i> forest on granite	B	Yog	Restricted to deep valleys and wet, fire-protected gullies on granite and Mathinna beds, often forming a sharp boundary with <i>E. sieberi</i> dry sclerophyll forest.
		Substrate not granite	SO	<i>E. sieberi</i> forest on other substrates	B	Yog	
WET-URN2	<i>E. urnigera</i> wsf		C	<i>E. coccifera</i> forest	A	N	Recorded only from shaded upper slopes of Mt Maria.

**Forest Botany Manual: Module 4 – Freycinet Region**

Floristic community code and name		Qualification	RFA community code and name		Conservation priority		Distribution in Freycinet Region
					Floristic	RFA	
WET-VIM0011	<i>E. viminalis</i> - <i>Bedfordia salicina</i> - <i>Pultenaea juniperina</i> wsf	<i>E. amygdalina</i> or <i>E. obliqua</i> codominant or subdominant	DSC	<i>E. viminalis</i> - <i>E. ovata</i> - <i>E. amygdalina</i> - <i>E. obliqua</i> damp sclerophyll forest	A	Yog	Wet sclerophyll - dry sclerophyll transition community found locally on dolerite slopes in the Eastern Tiers.
		Not DSC	VW	<i>E. viminalis</i> wet forest	A	Y*	
WET-VIM0100	<i>E. viminalis</i> - <i>Acacia dealbata</i> - <i>Pomaderris apetala</i> wsf		VW	<i>E. viminalis</i> wet forest	A	Y*	Local along creeklines and gullies in the Eastern Tiers, and also occurs as remnant forest along river flats (e.g. Buxton River).
WET-VIM0101	<i>E. viminalis</i> - <i>Acacia dealbata</i> - <i>Dicksonia antarctica</i> wsf		VW	<i>E. viminalis</i> wet forest	A	Y*	Local along humid creeklines and gullies in the Eastern Tiers (e.g. Anglers Creek).
WET-VIM2	<i>E. viminalis</i> - <i>Leptospermum lanigerum</i> - <i>Melaleuca squarrosa</i> wsf		OV	Shrubby <i>E. ovata</i> - <i>E. viminalis</i> forest	A	Y*	Very localised mainly on better-drained sites adjacent to flats with <i>E. ovata</i> forest, in higher rainfall parts of the region.

## 2.6 DRY SCLEROPHYLL FOREST AND WOODLAND COMMUNITIES

*Major Reference:* Duncan and Brown (1985)

Dry sclerophyll forests and woodlands are typically dominated by eucalypts under 40 m in height, and have a multi-layered understorey dominated by hard-leaved shrubs, including eucalypt regeneration. The eucalypts often form mixed species stands, and generally several age classes of eucalypts are present. The ground layer varies, but bracken, grasses and graminoids (sedges) are typical components. Many grassy communities are described in Kirkpatrick, Gilfedder and Fensham (1988).

Dry sclerophyll forest communities are classified by their overstorey dominants and the broad characteristics of their understorey. The understorey types are grassy (g), sedgy (sd), scrubby (sc), shrubby (sh) and heathy (h); they are described below. Inevitably, many areas will support vegetation with intermediate understoreys. FPOs need to use a precautionary approach when determining the conservation priority of such forests.

Land use practices (e.g. frequent or recent firing) can modify the structure or composition of dry sclerophyll understoreys. For example, frequent firing can lead to bracken displacing a diverse heathy understorey. FPOs should allow for land use practices when determining floristic communities.

Freycinet Region supports a greater range of dry sclerophyll forest and woodland communities than any other Tasmanian bioregion. Some dry sclerophyll communities, and many dry sclerophyll species, are restricted to Freycinet Region. Dry sclerophyll forests and woodlands are the major vegetation type in drier parts of the region (e.g. eastern Midlands and the central East Coast) and on infertile coastal sites (e.g. Freycinet Peninsula). At higher altitudes (e.g. Eastern Tiers) and more humid parts of the region (e.g. Wielangta), dry sclerophyll forests occur extensively on drier, more exposed sites, often with a high rock cover. Dry sclerophyll forest grades into wet sclerophyll forest (often through a damp sclerophyll transition zone) as sites become progressively more humid and less frequently burnt. As the environment becomes drier or more limiting to tree growth, dry sclerophyll forests grade into woodland, scrub, heath, grassland or sedgeland. Some dry sclerophyll communities in Freycinet Region, mainly associated with arable sites, have been extensively cleared or modified and have a high priority for conservation.

There are difficulties with allocating some dry sclerophyll floristic communities to RFA communities. These problem communities are not encountered in many FPP areas. The tables indicate when FPOs need to check with FPA for guidance on community affinities, priorities and prescriptions.

Many lowland dry sclerophyll communities in Freycinet Region, particularly on siliceous substrates and poorly drained sites, are highly susceptible to *Phytophthora cinnamomi*. These are indicated in the tables. Specialist advice will be needed if prescriptions in *Flora Technical Note 8* cannot be applied to operations in these communities.

### RECOGNISING THE DIFFERENT DRY SCLEROPHYLL UNDERSTOREYS

Each understorey type is recognised by the dominance or prominence of a distinctive suite of species. Species from other understorey types may also be present, and these communities will grade into one another in some situations, so it is important to note which species are the most dominant, rather than just which species are present. Eucalypt regrowth can be present in all understorey types.

#### Grassy forests

Grasses or sags are the dominant or most prominent feature of the understorey. Typical species include tussock grass, kangaroo grass, wallaby grass and sagg (*Lomandra longifolia*). Note that buttongrass and cutting grass are actually sedges, and forests with understoreys dominated by these species should be considered as sedgy communities. The ground layer generally contains a high diversity of herbs, most evident when they are flowering in spring and summer. Small trees and shrubs (e.g. black wattle, she-oak, prickly box) are widespread on drier lowland sites. The eucalypt canopy is often fairly open. Common species in Freycinet Region include *E. amygdalina*, *E. pulchella*, *E. globulus* and *E. viminalis*. Grassy forests are often found on relatively dry and well-drained sites on fertile substrates (e.g. dolerite), and flats subject to frost and cold air drainage.

### Sedgy forests

Sedges or rushes are the dominant or most prominent feature of the understorey. Typical species include cutting sedge, sword sedge, buttongrass, cutting grass and rushes. Coral ferns are often present. Shrubs such as tea-trees and paperbarks are present on many sites. The eucalypt canopy is fairly open. Common species in Freycinet Region include *E. amygdalina*, *E. ovata* and *E. rodwayi*. Sedgy forests grade into scrubby forests as shrub cover increases (in the absence of fire) and sedges reduce in cover. Sedgy forests occur on sites with impeded drainage, often on sites that have been burnt frequently or recently.

### Scrubby forests

Shrubs (typically tea-trees and paperbarks) are the dominant or most prominent feature of the understorey. They form a moderately dense to dense cover, generally over a sedgy ground layer. Other common shrub species include prickly mo, banksia, hakea and a range of legumes and heath species. The ground layer contains species typical of sedgy forest, though it is generally sparser, particularly under a dense shrub layer. The eucalypt canopy is typically fairly open. Common species in Freycinet Region include *E. amygdalina* and *E. ovata*. Scrubby forests mainly occur on flats with impeded drainage, generally on sites that have not been burnt or severely disturbed for many years. Scrubby forests often intergrade or form a mosaic with sedgy communities.

### Heathy forests

Shrubs less than 2 m in height are the dominant or most prominent feature of the understorey, though in frequently burnt sites this shrub layer can be displaced by bracken. Occasional taller shrubs are also often present in heathy forests. Shrub species include many heaths (e.g. *Epacris* species), legumes, wattles, bull-oak, banksia, tea-trees and grasstree (*Xanthorrhoea australis*). Bracken is the most widespread ground layer species, but sagg, sedges and colourful herbs (e.g. orchids, lilies) are often conspicuous. Eucalypt height and density varies in response to site conditions, but common species in Freycinet Region include *E. amygdalina* and *E. tenuiramis*. Heathy forests are generally found on well drained sites on infertile or siliceous substrates (e.g. sands, sandstone, quartzite, granite).

### Shrubby forests

Shrubs more than 2 m in height are the dominant or most prominent feature of the understorey. Several shrub layers are often present, often containing a mixture of wet sclerophyll (broad-leaved) and dry sclerophyll (narrow-leaved) shrubs. Shrubby forests are synonymous with damp sclerophyll forests when wet and dry sclerophyll shrubs are present in similar proportions. Common shrub species include native cherry, wattles, blanket bush, dolly bush, banksia, hop bush, prickly beauty, guitar plant and hakea. Ground layer species include bracken and other ferns, flax lily, sagg and grasses, though their cover is often sparse. Eucalypts are typically taller and denser than in other dry sclerophyll forest communities. In Freycinet Region they include *E. obliqua*, *E. delegatensis* and *E. globulus* on moist sites and *E. amygdalina* on drier sites. Shrubby forests tend to occupy more fertile sites, or more shaded and humid environments, than other dry sclerophyll types.

## KEY TO DRY SCLEROPHYLL FOREST AND WOODLAND COMMUNITIES

*Note: Exclude eucalypt regeneration in assessing dominance of understorey layers.*

- ① Understorey dominated by grasses and sagg; small trees or shrubs generally sparse; mainly on drier sites on basalt, dolerite or other fertile substrates
  - ② *E. amygdalina* dominant ..... DRY-gAM
  - ② *E. dalrympleana* dominant ..... DRY-gDAL
  - ② *E. delegatensis* dominant ..... DRY-gDEL
  - ② *E. globulus* dominant ..... DRY-gGLOB
  - ② *E. gunnii* dominant ..... DRY-gGUN
  - ② *E. ovata* dominant ..... DRY-gOV
  - ② *E. pauciflora* dominant ..... DRY-gPAUC
  - ② *E. pulchella* dominant ..... DRY-gPUL
  - ② *E. rodwayi* dominant ..... DRY-gROD
  - ② *E. rubida* dominant ..... DRY-gRUB
  - ② *E. tenuiramis* dominant ..... DRY-gTEN
  - ② *E. viminalis* dominant ..... DRY-gVIM

- ❶ **Understorey dominated by sedges, cutting grass or buttongrass or tea-trees or paperbarks; mainly on sites with impeded drainage (e.g. flats and marsh edges)**
- ❷ Sedges, cutting grass or buttongrass prominent
- ❸ *E. amygdalina* dominant ..... DRY-sdAM
  - ❸ *E. gunnii* dominant ..... DRY-sdGUN
  - ❸ *E. ovata* dominant ..... DRY-sdOV
  - ❸ *E. rodwayi* dominant ..... DRY-sdROD
- ❷ Tea-trees or paperbarks prominent
- ❸ *E. amygdalina* dominant ..... DRY-scAM
  - ❸ *E. gunnii* dominant ..... DRY-scGUN
  - ❸ *E. ovata* dominant ..... DRY-scOV
  - ❸ *E. rodwayi* dominant ..... DRY-scROD
- ❶ **Understorey dominated by bracken or low shrubs (generally less than 2 m), notably heaths, legumes, wattles, tea-trees, bull-oak and banksia; mainly on well drained sites on granite, sands, sandstone and other sediments**
- ❷ *E. amygdalina* dominant ..... DRY-hAM
  - ❷ *E. globulus* dominant ..... DRY-hGLOB
  - ❷ *E. pulchella* dominant ..... DRY-gPUL or DRY-shPUL
  - ❷ *E. obliqua* dominant ..... DRY-hOB
  - ❷ *E. ovata* dominant ..... DRY-hOV
  - ❷ *E. sieberi* dominant ..... DRY-hSIEB
  - ❷ *E. tenuiramis* dominant ..... DRY-hTEN
  - ❷ *E. viminalis* dominant ..... DRY-hVIM
- ❶ **Understorey dominated by shrubs over 2 m (excluding tea-trees and paperbarks), often including broad-leaved species; bracken or other ferns sometimes dense; mainly on well drained or sheltered sites of moderate fertility (e.g. dolerite)**
- ❷ *E. amygdalina* dominant ..... DRY-shAM
  - ❷ *E. cordata* dominant ..... DRY-shCORD
  - ❷ *E. dalrympleana* dominant ..... DRY-shDAL
  - ❷ *E. delegatensis* dominant ..... DRY-shDEL
  - ❷ *E. globulus* dominant ..... DRY-shGLOB
  - ❷ *E. obliqua* dominant ..... DRY-shOB
  - ❷ *E. ovata* dominant ..... DRY-shOV
  - ❷ *E. pauciflora* dominant ..... DRY-shPAUC
  - ❷ *E. pulchella* dominant ..... DRY-shPUL
  - ❷ *E. sieberi* dominant ..... DRY-shSIEB
  - ❷ *E. tenuiramis* dominant ..... DRY-shTEN
  - ❷ *E. viminalis* dominant ..... DRY-shVIM

**CONSERVATION PRIORITIES AND ATTRIBUTES OF DRY SCLEROPHYLL FOREST / WOODLAND COMMUNITIES**

Notes: # – Community highly susceptible to *Phytophthora cinnamomi* – specialist advice needed if prescriptions in *Flora Technical Note 8* cannot be applied

\* – Community identified as Rare, Vulnerable or Endangered at a Statewide level through RFA processes

Floristic community code and name		Qualification		RFA community code and name		Cons. priority		Distribution in Freycinet Region
						Floristic	RFA	
DRY-gAM	Grassy <i>E. amygdalina</i> forest/woodland	Substrate sand, alluvium, Tertiary gravels or ironstone (non-coastal areas)		AIC	Inland <i>E. amygdalina</i> forest on Cainozoic deposits	A	Y*	Local on broad flats in Cranbrook area and St Pauls Valley and Fingal Valley.
		Substrate dolerite (or basalt)	Not associated with AIC	AD	<i>E. amygdalina</i> forest on dolerite	np	N	Widespread, especially on the inland margins of the Eastern Tiers.
			Occurs with AIC	check	check	check	check	May occur locally in St Pauls and Fingal Valleys.
		Substrate mudstone		AM	<i>E. amygdalina</i> forest on mudstone	check	check	May occur locally on dry sites in SE of region.
		Substrate mudstone		check	check	check	check	
DRY-gDAL	Grassy <i>E. dalrympleana</i> forest/woodland	Make sure dominant is not <i>E. viminalis</i> - contact FPA if unsure.		D	Dry <i>E. delegatensis</i> forest	B	N	Local on well drained slopes and flats at higher altitudes (e.g. Lake Leake area).
DRY-gDEL	Grassy <i>E. delegatensis</i> forest/woodland			D	Dry <i>E. delegatensis</i> forest	np	N	Cool, dry upland dolerite slopes and tier surfaces (e.g. Mt Foster).
DRY-gGLOB	Grassy <i>E. globulus</i> forest/woodland	<i>E. pulchella</i> present as codominant		P	<i>E. pulchella</i> - <i>E. globulus</i> - <i>E. viminalis</i> grassy shrubby forest	np	N	Dolerite slopes and ridges, particularly in the Eastern Tiers and its coastal foothills (e.g. Mayfield, Lisdillon Hills).
		<i>E. pulchella</i> present as subdominant or minor species or absent		GG	Grassy <i>E. globulus</i> forest	A	Y*	Mainly on dolerite in coastal areas (e.g. Bangor, Earlham, Ryton Hills) and eastern Midlands (e.g. Stonehenge). Also local on granite (e.g. Four Mile Creek) and sandstone (e.g. Sawpit Tier).
DRY-gGUN	Grassy <i>E. gunnii</i> forest/woodland			C	<i>E. coccifera</i> forest	A	N	Very localised on high altitude tiers subject to frost and poor drainage (e.g. Snow Hill, Mt Foster).
DRY-gOV	Grassy <i>E. ovata</i> forest/woodland			OV	Shrubby <i>E. ovata</i> - <i>E. viminalis</i> forest	A <sup>#</sup>	Y*	Local on low altitude flats with impeded drainage, often frequently fired (e.g. Cusicks Hill area).
DRY-gPAUC	Grassy <i>E. pauciflora</i> forest/woodland			PJ	<i>E. pauciflora</i> forest on Jurassic dolerite	B	Yog	Well drained frost hollows and flats, mainly on dolerite (e.g. Lake Leake area, Thompsons Marsh).
DRY-gPUL	Grassy <i>E. pulchella</i> forest/woodland	May occur locally with a heathy understorey		P	<i>E. pulchella</i> - <i>E. globulus</i> - <i>E. viminalis</i> grassy shrubby forest	np	N	Widespread on dry dolerite slopes and ridges, particularly in the Eastern Tiers and its coastal foothills.

**Forest Botany Manual: Module 4 – Freycinet Region**

Floristic community code and name		Qualification	RFA community code and name		Cons. priority		Distribution in Freycinet Region
					Floristic	RFA	
DRY-gROD	Grassy <i>E. rodwayi</i> forest/woodland		RO	<i>E. rodwayi</i> forest	A	Y	Local on poorly drained flats, often frequently fired, in uplands and frost hollows in eastern Midlands (e.g. Ladies Mile Marsh, Ginger Marsh).
DRY-gRUB	Grassy <i>E. rubida</i> forest/woodland		V	<i>E. viminalis</i> grassy forest/woodland	A	Y	Local on dolerite flats, saddles and frost hollows in the eastern Midlands.
DRY-gTEN	Grassy <i>E. tenuiramis</i> forest/woodland	Substrate dolerite	TD	<i>E. tenuiramis</i> forest on dolerite	np	N	Local on well drained dolerite flats and broad ridges, particularly on Tasman Peninsula.
		Substrate mudstone or sandstone	TI	Inland <i>E. tenuiramis</i> forest	A	Y*	Mainly found on dry mudstone sites, often frequently burnt (e.g. Mt Elizabeth, Bangor).
DRY-gVIM	Grassy <i>E. viminalis</i> forest/woodland	Substrate sand, alluvium, Tertiary gravels or ironstone	AIC	Inland <i>E. amyg.</i> forest on Cainozoic deposits	A	Y*	Localised on broad flats in Cranbrook area, St Pauls Valley and Fingal Valley.
		Substrate dolerite (or basalt)	V	<i>E. viminalis</i> grassy forest/woodland	A	Y	Common on dry dolerite hills and slopes of eastern Midlands (e.g. Pepper Creek), Eastern Tiers (e.g. Grange Hills, Belmont) and Maria Island.
		Substrate other (mainly mudstone or Mathinna series)	check	check	check	check	Occasional, mainly on sediments. Generally associated with <i>E. amygdalina</i> forest.
DRY-hAM	Heathy <i>E. amygdalina</i> forest	Substrate sand or alluvium in (sub)coastal areas or granite	AC	Coastal <i>E. amygdalina</i> forest	np <sup>#</sup>	N	Widespread on sand and granite in coastal areas (e.g. Freycinet NP, Richardsons Road); extending inland on some sites (e.g. Royal George).
		Substrate sandstone (mainly Triassic, also Mathinna series)	AS	<i>E. amygdalina</i> forest on sandstone	A <sup>#</sup>	Y*	Well drained sandstone sites (e.g. Buckland area, Mt Douglas).
		Substrate sand, alluvium, Tertiary gravels or ironstone (non-coastal areas)	AIC	Inland <i>E. amygdalina</i> forest on Cainozoic deposits	A	Y*	Local on broad flats in Cranbrook area, St Pauls Valley and Fingal Valley.
		Substrate mudstone or Mathinna series	AM	<i>E. amygdalina</i> forest on mudstone	check	check	Mainly dry slopes on Mathinna sediments in Mt Foster and Dog Kennels area.
DRY-hGLOB	Heathy <i>E. globulus</i> forest		G	<i>E. viminalis</i> and/or <i>E. globulus</i> coastal shrubby forest	A <sup>#</sup>	Y*	Local on well drained sites on Recent sand in coastal areas (e.g. McRaes Isthmus (Maria Island), Rheban Spit, Crescent Bay).
DRY-hOB	Heathy <i>E. obliqua</i> forest		O	<i>Dry E. obliqua</i> forest	np <sup>#</sup>	N	Local on granite, sandstone or sand in coastal and inland areas (e.g. Buckland, Tasman Peninsula).

**Forest Botany Manual: Module 4 – Freycinet Region**

Floristic community code and name		Qualification	RFA community code and name		Cons. priority		Distribution in Freycinet Region
					Floristic	RFA	
DRY-hOV	Heathy <i>E. ovata</i> forest/woodland		OV	Shrubby <i>E. ovata</i> - <i>E. viminalis</i> forest	A <sup>#</sup>	Y*	Local on poorly drained flats, mainly in coastal areas (e.g. Freycinet Peninsula).
DRY-hSIEB	Heathy <i>E. sieberi</i> forest		SG	<i>E. sieberi</i> forest on granite	np <sup>#</sup>	Yog	Local on granite, in (sub)coastal areas in north of region (e.g. Friendly Beaches, Bedgood Hill).
DRY-hTEN	Heathy <i>E. tenuiramis</i> forest	Substrate granite	TG	<i>E. tenuiramis</i> forest on granite	np <sup>#</sup>	N	Local on well drained granite sites in Freycinet area (e.g. Cape Tourville).
		Substrate sand	check	check	check	check	Local on sand, often wind-blown (e.g. Taranna).
		Substrate sandstone or mudstone	TI	Inland <i>E. tenuiramis</i> forest	A <sup>#</sup>	Y*	Local on well drained sites on mudstone or sandstone (e.g. Mt Douglas area).
DRY-hVIM	Heathy <i>E. viminalis</i> forest	Substrate coastal sand	G	<i>E. viminalis</i> and/or <i>E. globulus</i> coastal shrubby forest	A <sup>#</sup>	Y*	Local on well drained sites on Recent sand in coastal areas (e.g. McRaes Isthmus (Maria Island), Rheban Spit, Crescent Bay, Dolphin Sands).
		Substrate granite	AC	Coastal <i>E. amygdalina</i> forest	A <sup>#</sup>	N	May occur locally on granite in coastal and subcoastal areas (e.g. Friendly Beaches).
		Substrate sand or alluvium (non-coastal areas) or Tertiary ironstone or gravels	AIC	Inland <i>E. amygdalina</i> forest on Cainozoic deposits	A	Y*	May occur locally on broad valley flats in inland areas (e.g. Cranbrook area, St Pauls Valley and Fingal Valley).
		Substrate other (mainly mudstone or Mathinna series)	check	check	check	check	Occasional on other substrates (e.g. sandstone, mudstone), generally with <i>E. amygdalina</i> forest.
DRY-sdAM <u>and</u> DRY-scAM	Sedgy <i>E. amygdalina</i> forest/woodland  Scrubby <i>E. amygdalina</i> forest/woodland	Substrate sand or alluvium in (sub)coastal areas <u>or</u> granite <u>or</u> Precambrian beds	AC	Coastal <i>E. amygdalina</i> forest	np <sup>#</sup>	N	Local around drainage lines and margins of marshes on a range of substrates (e.g. Wrights Marsh, Thompsons Marsh, Coal Marsh).
	Substrate sandstone (mainly Triassic and Ordovician)	AS	<i>E. amygdalina</i> forest on sandstone	A <sup>#</sup>	Y*		
	Substrate sand, alluvium, Tertiary gravels or ironstone (non-coastal areas)	AIC	Inland <i>E. amygdalina</i> forest on Cainozoic deposits	A	Y*		
	Substrate dolerite (or basalt)	Not associated with AIC	AD	<i>E. amygdalina</i> forest on dolerite	np	N	
		Occurs with AIC	check	check	check	check	
	Substrate other	check	check	check	check		

**Forest Botany Manual: Module 4 – Freycinet Region**

Floristic community code and name		Qualification	RFA community code and name		Cons. priority		Distribution in Freycinet Region
					Floristic	RFA	
DRY-sdGUN <u>and</u> DRY-scGUN	Sedgy <i>E. gunnii</i> forest/woodland Scrubby <i>E. gunnii</i> forest/woodland		C	<i>E. coccifera</i> forest	A	N	Very localised on poorly drained flats at high altitudes (e.g. Mt Foster, Snow Hill).
DRY-sdOV <u>and</u> DRY-scOV	Sedgy <i>E. ovata</i> forest/woodland Scrubby <i>E. ovata</i> forest/woodland		OV	Shrubby <i>E. ovata</i> - <i>E. viminalis</i> forest	A <sup>#</sup>	Y*	Widespread but local on poorly drained flats, particularly in lowland areas (e.g. Swilly Marsh).
DRY-sdROD <u>and</u> DRY-scROD	Sedgy <i>E. rodwayi</i> forest/woodland Scrubby <i>E. rodwayi</i> forest/woodland		RO	<i>E. rodwayi</i> forest	A	Y	Local on poorly drained flats, particularly in upland areas (e.g. Flagstaff Marsh) but also at lower altitudes (e.g. Dolans Marsh).
DRY-shAM	Shrubby <i>E. amygdalina</i> forest	Understorey with similar amount of wet and dry sclerophyll shrubs <u>and</u> <i>E. obliqua</i> or <i>E. viminalis</i> subdominant or codominant	DSC	<i>E. viminalis</i> - <i>E. ovata</i> - <i>E. amygdalina</i> - <i>E. obliqua</i> damp sclerophyll forest	np	Yog	Found on sites intermediate between wet and dry forest. Common on shaded slopes in drier parts of the region, and moderately exposed slopes in wetter parts of the region (e.g. Wielangta, Buckland Training Area, Eastern Tiers)
		Not DSC and substrate dolerite (or basalt)	AD	<i>E. amygdalina</i> forest on dolerite	np	N	Widespread on dolerite, particularly in the Eastern Tiers.
		Not DSC and substrate sand or granite	AC	Coastal <i>E. amygdalina</i> forest	np <sup>#</sup>	N	Local on moderately shaded slopes on granite and possibly sand.
		Not DSC and substrate sandstone	AS	<i>E. amygdalina</i> forest on sandstone	A <sup>#</sup>	Y*	Local on moderately shaded slopes on sandstone.
		Not DSC and substrate mudstone or Mathinna series	AS	<i>E. amygdalina</i> forest on sandstone	check	check	Local on moderately shaded slopes and gully flanks or long unburnt sites.
		Not DSC and substrate other	check	check	check	check	Local on moderately shaded slopes.
DRY-shCOC	Shrubby <i>E. coccifera</i> forest		C	<i>E. coccifera</i> forest	A	N	Local on exposed dolerite knolls and ridges at mid to high altitudes (e.g. Snow Hill, Middle Peak).
DRY-shCORD	Shrubby <i>E. cordata</i> forest		check	check	A	check	Very local, often on dolerite knolls and saddles, particularly in the southeast of the region (e.g. Wielangta area, Bluestone Tier).

**Forest Botany Manual: Module 4 – Freycinet Region**

Floristic community code and name		Qualification	RFA community code and name		Cons. priority		Distribution in Freycinet Region
					Floristic	RFA	
DRY-shDAL	Shrubby <i>E. dalrympleana</i> forest	Make sure dominant is not <i>E. viminalis</i> - contact FPA if unsure.	D	Dry <i>E. delegatensis</i> forest	B	N	Local on well drained slopes and flats at higher altitudes; often associated with DRY-shDEL.
DRY-shDEL	Shrubby <i>E. delegatensis</i> forest		D	Dry <i>E. delegatensis</i> forest	np	N	Widespread on moderately exposed slopes, ridges and tier surfaces at higher altitudes on rocky (usually dolerite), free draining sites (e.g. Mt Foster, Mt Tooms).
DRY-shGLOB	Shrubby <i>E. globulus</i> forest	<i>E. pulchella</i> present as codominant	P	<i>E. pulchella</i> - <i>E. globulus</i> - <i>E. viminalis</i> grassy shrubby forest	np	N	Common on shaded, rocky or infrequently burnt dolerite sites (e.g. Eastern Tiers, Wielangta, Mt Morrison area).
		<i>E. pulchella</i> present as subdominant or minor species or absent	GG	Grassy <i>E. globulus</i> forest	A	Y*	Mainly dolerite sites where grassy understorey is replaced by shrubbier understorey because of moister conditions or less frequent fire.
		Substrate sand or granite in coastal areas	G	<i>E. viminalis</i> and/or <i>E. globulus</i> coastal shrubby forest	A <sup>#</sup>	Y*	Local on humid or fire-free coastal sites that would normally support heathy <i>E. globulus</i> forest.
		Not P, GG or G	check	check	check	check	Local on humid or fire-free sites that would normally support more open <i>E. globulus</i> forest.
DRY-shOB	Shrubby <i>E. obliqua</i> forest	Understorey with similar amount of wet and dry sclerophyll shrubs <u>and</u> <i>E. amygdalina</i> or <i>E. viminalis</i> subdominant or codominant	DSC	<i>E. viminalis</i> - <i>E. ovata</i> - <i>E. amygdalina</i> - <i>E. obliqua</i> damp sclerophyll forest	np	Yog	Found on sites intermediate between wet and dry forest. Common on shaded slopes in drier parts of the region, and moderately exposed slopes in wetter parts of the region (e.g. Wielangta, Buckland Training Area, Eastern Tiers)
		Not DSC	O	Dry <i>E. obliqua</i> forest	np	N	Widespread in moister lowland environments on less exposed slopes and gully flanks.
DRY-shOV	Shrubby <i>E. ovata</i> forest		OV	Shrubby <i>E. ovata</i> - <i>E. viminalis</i> forest	A <sup>#</sup>	Y*	Local on sites intermediate between DRY-scOV and wet sclerophyll forest.
DRY-shPAUC	Shrubby <i>E. pauciflora</i> forest/woodland	Substrate dolerite (or basalt)	PJ	<i>E. pauciflora</i> forest on Jurassic dolerite	B	Yog	Local on dolerite flats in Eastern Tiers subject to frost and cold air drainage (e.g. Lake Leake area).
		Substrate other	check	check	check	check	Occasional, mainly on sediments (e.g. Buckland).
DRY-shPUL	Shrubby <i>E. pulchella</i> forest	May occur locally with a heathy understorey	P	<i>E. pulchella</i> - <i>E. globulus</i> - <i>E. viminalis</i> grassy shrubby forest	np	N	Common on partly shaded dolerite slopes, often forming a transition zone between wet forests and dry forests (e.g. Eastern Tiers, Wielangta).

**Forest Botany Manual: Module 4 – Freycinet Region**

Floristic community code and name		Qualification	RFA community code and name		Conservation priority		Distribution in Freycinet Region
					Floristic	RFA	
DRY-shSIEB	Shrubby <i>E. sieberi</i> forest	Substrate sand or granite	SG	<i>E. sieberi</i> forest on granite	np <sup>#</sup>	Yog	Local in north of region, mainly on granite (e.g. Wardlaws Creek area).
		Other substrates	SO	<i>E. sieberi</i> forest on other substrates	np <sup>#</sup>	Yog	Local in north of region on dolerite (e.g. Mt Allen) or Mathinna sediments (e.g. Elephant Pass area).
DRY-shTEN	Shrubby <i>E. tenuiramis</i> forest	Substrate granite	TG	<i>E. tenuiramis</i> forest on granite	np <sup>#</sup>	N	Local on well drained granite sites in Freycinet area (e.g. Cape Tourville).
		Substrate dolerite	TD	<i>E. tenuiramis</i> forest on dolerite	np	N	Local on lower slopes and flats, often near dolerite-sediment contact zones (e.g. Denison Marshes) and on Tasman Peninsula.
		Substrate sand	check	check	check	check	Local on sand (often wind-blown) e.g. Taranna.
		Substrate sandstone or mudstone	TI	Inland <i>E. tenuiramis</i> forest	A	Y*	Local on well drained sites on mudstone or sandstone (e.g. Mt Douglas area).
DRY-shVIM	Shrubby <i>E. viminalis</i> forest	Understorey with similar amount of wet and dry sclerophyll shrubs <u>and</u> <i>E. amygdalina</i> or <i>E. obliqua</i> subdominant or codominant	DSC	<i>E. viminalis</i> - <i>E. ovata</i> - <i>E. amygdalina</i> - <i>E. obliqua</i> - damp sclerophyll forest	A	Yog	Local in dry sclerophyll - wet sclerophyll transition zones, sometimes along creeklines in drier areas.
		Not DSC and substrate dolerite (or basalt)	V	<i>E. viminalis</i> grassy forest/woodland	A	Y	Local on humid or fire-free coastal sites that would normally support grassy <i>E. viminalis</i> forest.
		Not DSC and substrate coastal sand	G	<i>E. viminalis</i> and/or <i>E. globulus</i> coastal shrubby forest	A <sup>#</sup>	Y*	Local on humid or fire-free coastal sites that would normally support heathy <i>E. viminalis</i> forest.
		Not DSC and substrate granite or (sub)coastal gravels	AC	Coastal <i>E. amygdalina</i> forest	B <sup>#</sup>	N	Occasional, usually with <i>E. amygdalina</i> forest.
		Not DSC and substrate other	check	check	check	check	Occasional, usually with <i>E. amygdalina</i> forest.

## Section 3 PRIORITY SPECIES

Species listed in this section have some priority for conservation, and are known or likely to occur in Freycinet Region. Most are listed on Schedules of the Tasmanian *Threatened Species Protection Act* 1995, with a small proportion also being listed under the Commonwealth *Environment Protection and Biodiversity Conservation Act* 1999. This section also includes a few species (e.g. outliers of biogeographic interest) that are not listed under either Act. Priority species are arranged on the basis of the timber quality of the forests that they typically occupy, but may also occur in other vegetation types. Some priority species have been recorded from plantations.

Much more information is available on the taxonomy, distribution and conservation status of vascular species (ferns, conifers and flowering plants) than non-vascular species (mosses, liverworts and lichens). This is partly because more texts are available on vascular plants, and partly because identification of non-vascular species often requires microscopic examination of specimens.

Our knowledge of the distribution and ecology of threatened species is improving as new information is obtained from targeted studies and from surveys in different areas of Tasmania (including surveys conducted by forest workers).. Information from all sources allows the conservation status and requirements of listed species to be reviewed and better prescriptions to be developed.

In many cases, assessments can be directed towards particular environments (e.g. dry rocky knolls) because many threatened species have narrow habitat ranges, and potential occurrences can be predicted from conditions of the physical environment (e.g. geology, aspect, altitude) or biological environment (e.g. dominant tree species, plant community). Many sites containing listed species will be flagged as priority communities (Section 2) or sites of potential significance for flora (Section 4).

Many threatened species are known or likely (based on taxonomic affinities) to be susceptible to *Phytophthora cinnamomi*. These species are indicated in the tables of priority species. Many of these species are found on lowland, siliceous sites. Some of the species play an important role in ecosystem functioning. Particular care is needed in evaluating FPP areas and planning operations in environments that contain *Phytophthora*-susceptible species (see Section 6 and *Flora Technical Note 8*).

The FPP flora evaluation requires that FPOs use databases to determine if threatened species have been recorded from within or close to proposed operational areas. The nominated databases are GTSpot (DPIWE) and NewCONSERVE (Forestry Tasmania). Details of how to access these databases are given in Module 1. Both databases provide other information that may be useful in preparing FPPs. The databases are updated regularly to incorporate new records of threatened species. Updated information on threatened species in this bioregion may also be found on the FPA Website.

The absence of records does not mean that threatened species are not present – many FPP areas will not have been surveyed previously. If new sites containing threatened species are found, details on site location, abundance of the species, and other potentially useful information (e.g. habitat, land use and fire history) should be forwarded to FPA. Material (e.g. photos or scans) can also be sent to the FPA Botany or Ecology programs for confirmation.

Many priority species will not be familiar to FPOs, but some are readily identifiable. Information to help identify many of the listed species is available on the FPA Website and in some of the references in *Flora Technical Note 2*. Further information on threatened plant species can be obtained from the Threatened Flora of Tasmania website (<http://www.gisparks.tas.gov.au/ThreatenedFloraCD/>). This site contains individual PDF files of all plant species listed on Schedules of the *Threatened Species Protection Act*. The files contain an image of the species (which could be useful in field verification) as well as information on habitat, distribution and conservation management.

Contact FPA if any priority species are identified or are likely to occur in an area that could be affected by a forestry operation. Recommendations to take account of such occurrences will be developed in conjunction with DPIWE (Threatened Species Unit). They will depend on characteristics of the species, site and proposed operation. Some sites may need to be excluded from operations, but often the values can be maintained by management prescriptions. In some cases (e.g. for opportunistic species) no changes to the proposed operation will be needed.

**PRIORITY SPECIES ARRANGED BY BROAD VEGETATION TYPES**

Priority species are arranged by broad vegetation type: corresponding PI typing is also indicated. More information on distribution and habitat can be obtained from the Threatened Flora of Tasmania website (<http://www.gisparks.tas.gov.au/ThreatenedFloraCD/>).

The status of the species refers to its presence on Schedules of the *Threatened Species Protection Act*:

- X Presumed extinct in Tasmania (Schedule 3)
- E Endangered in Tasmania (Schedule 3)
- V Vulnerable in Tasmania (Schedule 4)
- R Rare and at risk in Tasmania (Schedule 5)

Susceptibility of species (in their natural habitat) to *Phytophthora cinnamomi* (Pc) is indicated. This is based on Barker and Wardlaw (1995) and Schahinger *et al.* (2003):

- Hs Highly susceptible: expect >75% mortality of infected plants to be killed
- Ms Moderately susceptible: expect 25-75% mortality of infected plants
- Prb Probably highly or moderately susceptible but no records of *Phytophthora* infection
- Ss Slightly susceptible: symptomless but reduced vigour
- S Susceptible but unable to make a rating
- Rh Resistant host: *Phytophthora* persists but host shows no symptoms.

**FOREST QUALITY MODERATE TO HIGH (e.g. E3+, E2, M+)**

Life form	Status	Pc	Botanical name	Common name
Trees & shrubs	R	Ss	<i>Acacia pataczekii</i>	Wally's wattle
		Prb	<i>Cyathodes platystoma</i>	Giant cheeseberry
	R		<i>Gynatrix pulchella</i>	Hemp bush
	R	Ms	<i>Hovea corrickiae</i>	Glossy hovea
		Hs	<i>Phyllocladus aspleniifolius</i>	Celery-top pine
Ferns	V		<i>Asplenium hookerianum</i>	Hooker's spleenwort
	V		<i>Blechnum cartilagineum</i>	Gristle fern
	E		<i>Cyathea cunninghamii</i>	Slender tree fern
	V		<i>Cyathea Xmarcescens</i>	Skirted tree fern
	R		<i>Hypolepis muelleri</i>	Harsh ground fern
Herbs	R		<i>Austrocynoglossum latifolium</i>	Forest hound's tongue
	R		<i>Plantago debilis</i>	Shade plantain
	R		<i>Senecio velleioides</i>	Forest groundsel
Orchids			<i>Sarcochilus australis</i>	Gunn's tree orchid
Grasses	R		<i>Deyeuxia benthamiana</i>	Bentham's bent grass
	R		<i>Hierochloe rariflora</i>	Cane holy grass
Other monocots	R		<i>Carex gunniana</i>	Mountain sedge

**FOREST QUALITY LOW TO MODERATE (e.g. E4, E3-)**

Note that many species listed above for moderate to high quality forests extend to lower quality forests.

Life form	Status	Pc	Botanical name	Common name
Trees & shrubs	V	Ms	<i>Acacia axillaris</i>	Midlands wattle
	R	Hs	<i>Acacia mucronata</i> subsp. <i>dependens</i>	Variable sawtooth wattle
	R	Hs	<i>Acacia ulicifolia</i>	Juniper wattle
	V		<i>Bertya tasmanica</i> subsp. <i>tasmanica</i>	Bertya
	V	Prb	<i>Boronia hippopala</i>	Velvet boronia
	R	Ss	<i>Bossiaea obcordata</i>	Spiny bossiaea
	R		<i>Cyphanthera tasmanica</i>	Tasmanian ray flower
	R	Ss	<i>Epacris acuminata</i>	Clasping-leaf heath
	E		<i>Epacris apseyensis</i>	Apsley heath
	V	Ms	<i>Epacris exserta</i>	South Esk heath
	V	Hs	<i>Epacris grandis</i>	Great heath
	E	Hs	<i>Epacris limbata</i>	Border heath

**Forest Botany Manual: Module 4 – Freycinet Region**

Life form	Status	Pc	Botanical name	Common name
	V	Ms	<i>Epacris virgata</i> aff. <i>Kettering</i>	Southern pretty heath
<b>Trees &amp; shrubs</b>			<i>Eucalyptus archeri</i>	Alpine cider gum
			<i>Eucalyptus cordata</i>	Heart-leaf silver gum
			<i>Eucalyptus urnigera</i>	Urn gum
	R		<i>Grevillea australis</i> var. <i>linearifolia</i>	Narrow-leaf southern grevillea
	R		<i>Odixia achlaena</i>	Odixia
	R		<i>Olearia hookeri</i>	Hooker's crimson tip daisy bush
	R		<i>Ozothamnus lycopodioides</i>	Lycopoid everlasting
	R	Prb	<i>Pentachondra ericifolia</i>	Matted pentachondra
	R		<i>Pimelea flava</i> subsp. <i>flava</i>	Yellow rice flower
	V		<i>Polyscias</i> aff. <i>sambucifolia</i>	Elderberry panax
	V		<i>Pomaderris elachophylla</i>	Small leaf pomaderris
	R		<i>Pomaderris intermedia</i>	Tree pomaderris
	R		<i>Pomaderris phyllicifolia</i> ssp. <i>phyllicifolia</i>	Narrow leaf pomaderris
	V		<i>Spyridium lawrencei</i>	Small leaf spyridium
	R		<i>Spyridium vexilliferum</i>	Winged spyridium
	V		<i>Stenanthemum pimeleoides</i>	Spreading stenanthemum
	R	S	<i>Thryptomene micrantha</i>	Heath myrtle, ribbed thryptomene
<b>Herbs</b>	R		<i>Brachyscome radicata</i>	Rooted daisy
	R		<i>Brachyscome sieberi</i> var. <i>gunnii</i>	Sieber's daisy
	V		<i>Brunonia australis</i>	Blue pincushion
	R		<i>Colobanthus curtisiae</i>	Curtis' colobanth
	R		<i>Cynoglossum australe</i>	Australian hound's tongue
	V		<i>Desmodium gunnii</i>	Slender or variable tick trefoil
	V		<i>Eryngium ovinum</i>	Blue devil
	V		<i>Glycine latrobeana</i>	Dwarf, purple or clover glycine
	V		<i>Glycine microphylla</i>	Small-leaf glycine
	R		<i>Lobelia rhombifolia</i>	Branched lobelia
	R		<i>Stellaria multiflora</i>	Rayless starwort
	R		<i>Teucrium corymbosum</i>	Forest germander
	R		<i>Veronica plebeia</i>	Trailing speedwell
<b>Orchids</b>	E		<i>Caladenia anthracina</i>	Black-tipped spider orchid
	E		<i>Prasophyllum apoxychilum</i>	Tapered leek orchid
	E		<i>Pterostylis atriola</i>	Snug greenhood
	E		<i>Pterostylis cycnocephala</i>	Swan greenhood
	R		<i>Pterostylis grandiflora</i>	Superb or cobra greenhood
	R		<i>Pterostylis squamata</i>	Ruddy greenhood
<b>Grasses</b>	E		<i>Aristida benthamii</i>	Three-awned speargrass
	R		<i>Deyeuxia apsleyensis</i>	Apsley bent grass
	R		<i>Deyeuxia brachyathera</i>	Short bent grass
	R		<i>Deyeuxia decipiens</i>	Bent grass
	R		<i>Rytidosperma procerum</i>	Tall wallaby grass
<b>Other monocots</b>	R		<i>Aphelia gracilis</i>	Slender aphelia
	R		<i>Arthropodium strictum</i>	Chocolate lily
	R		<i>Uncinia elegans</i>	Handsome hook sedge

**FOREST QUALITY VERY LOW OR NON-FOREST (e.g. E5, S, Vz)**

Note that many species listed above for low quality forests extend to very low quality forest or non-forest vegetation. Some species listed below have also been recorded within or adjacent to FPP areas.

Life form	Status	Pc	Botanical name	Common name
<b>Trees &amp; shrubs</b>	R	Ms	<i>Acacia siculiformis</i>	Dagger wattle
	R		<i>Allocasuarina crassa</i>	Capes she-oak
	V	Prb	<i>Boronia gunnii</i>	Gunn's boronia
	R		<i>Brachyloma depressum</i>	Spreading brachyloma
	V		<i>Callitris oblonga</i> subsp. <i>oblonga</i>	South Esk pine
	V	S	<i>Conospermum hookeri</i>	Variable smoke bush
	E		<i>Cryptandra amara</i>	Bitter cryptandra
			<i>Cryptandra exilis</i>	Coastal cryptandra
	E		<i>Discaria pubescens</i>	Thorn bush, anchor plant
	E	Hs	<i>Epacris barbata</i>	Bearded heath

Forest Botany Manual: Module 4 – Freycinet Region

		Hs	<i>Epacris marginata</i>	Stiffly erect heath
		R	<i>Eucalyptus barberi</i>	Barber's gum
Life form	Status	Pc	Botanical name	Common name
Trees & shrubs			<i>Eucalyptus coccifera</i>	Snow peppermint, Snow gum
	R		<i>Eutaxia microphylla</i>	Common eutaxia
	R		<i>Gyrostemon thesioides</i>	Broom wheel fruit
	E		<i>Hardenbergia violacea</i>	Happy wanderer
	R		<i>Hovea tasmanica</i>	Tasman hovea
	V		<i>Lasiopetalum micranthum</i>	Tasmanian velvet bush
	R		<i>Leucopogon lanceolatus</i>	Lance beard heath
	R		<i>Melaleuca pustulata</i>	Dry shrubby paperbark
	R	S	<i>Monotoca submutica</i> var. <i>autumnalis</i>	Round leaf broom heath
	R		<i>Muehlenbeckia axillaris</i>	Matted lignum
	E	Prb	<i>Philotheca freyciana</i>	Freycinet wax flower
	R		<i>Pomaderris oraria</i>	Coast pomaderris
	V	Rh	<i>Prostanthera rotundifolia</i>	Round leaf mint bush
	V	Hs	<i>Pultenaea prostrata</i>	Bush pea
	E		<i>Spyridium eriocephalum</i>	Heath spyridium
	R		<i>Spyridium parvifolium</i> var. <i>parvifolium</i>	Australian dusty miller
	V		<i>Stonesiella selaginoides</i>	Clubmoss bush pea
	E		<i>Viminaria juncea</i>	Golden spray
	R		<i>Wilsonia humilis</i>	Silky wilsonia
	R		<i>Wilsonia rotundifolia</i>	Round leaf wilsonia
	R		<i>Zieria littoralis</i>	Downy or dwarf zieria
	R		<i>Zieria veronicea</i> subsp. <i>veronicea</i>	Pink zieria
Ferns	R		<i>Cheilanthes distans</i>	Bristly cloak fern
	E		<i>Isoetes drummondii</i> subsp. <i>drummondii</i>	Plain quillwort
	R		<i>Isoetes elatior</i>	Tall quillwort
	R		<i>Pellaea calidirupium</i>	Hot rock fern
	R		<i>Pilularia novae-hollandiae</i>	Austral pillwort
			<i>Schizaea asperula</i>	Comb fern
Herbs	R		<i>Asperula subsimplex</i>	Water woodruff
	V		<i>Brachyscome rigidula</i>	Hairy cut-leaf daisy
	R		<i>Calocephalus lacteus</i>	Milky beauty heads
	R		<i>Calystegia soldanella</i>	Sea bindweed
	R	Prb	<i>Comesperma defoliatum</i>	Leafless milkwort
	R		<i>Cuscuta tasmanica</i>	Golden dodder
	R		<i>Euphrasia amphisepala</i>	Forestier eyebright
	R		<i>Euphrasia collina</i> subsp. <i>deflexifolia</i>	Eastern eyebright
	R		<i>Euphrasia collina</i> subsp. <i>gunnii</i>	Gunn's purple eyebright
	V		<i>Euphrasia phragmostoma</i>	Buften's eyebright
	E		<i>Euphrasia scabra</i>	Yellow eyebright
	E		<i>Euphrasia semipicta</i>	Port Arthur eyebright
	E		<i>Euphrasia</i> sp. <i>fabula</i>	Masked cliff eyebright
	R		<i>Glossostigma elatinoides</i>	Small mudmat
	R		<i>Goodenia geniculata</i>	Bent goodenia
	V		<i>Haloragis aspera</i>	Rough raspwort
	R		<i>Haloragis heterophylla</i>	Variable raspwort
	R		<i>Haloragis myriocarpa</i>	Prickly raspwort
	E		<i>Hyalosperma demissum</i>	Moss sunray
	V		<i>Hydrocotyle laxiflora</i>	Stinking pennywort
	E		<i>Isoetopsis graminifolia</i>	Grass cushions
	E		<i>Lepidium hyssopifolium</i>	Peppergrass
	R		<i>Lepidium pseudotasmanicum</i>	Peppergrass
	R		<i>Limonium australe</i>	Sea lavender
	V		<i>Limonium baudinii</i>	Baudin's sea lavender
	V		<i>Lobelia pratioides</i>	Poison lobelia
	R		<i>Lotus australis</i>	Austral trefoil
	R		<i>Millotia muelleri</i>	Common bow flower
	V		<i>Myriophyllum integrifolium</i>	Tiny water milfoil
	V		<i>Persicaria decipiens</i>	Slender knotweed
	V		<i>Phyllangium divergens</i>	Wiry mitrewort
	R		<i>Ranunculus sessiliflorus</i>	Small flowered Australian buttercup
	R		<i>Rhodanthe anthemoides</i>	Chamomile sunray

Forest Botany Manual: Module 4 – Freycinet Region

Life form	Status	Pc	Botanical name	Common name
	E		<i>Scaevola aemula</i>	Fairy fan flower
	R		<i>Scleranthus brockiei</i>	Brock knawel
<b>Herbs</b>	V		<i>Scleranthus diander</i>	Knawel
	V		<i>Scleranthus fasciculatus</i>	Spreading knawl
	R		<i>Scutellaria humilis</i>	Dwarf scullcap
	R		<i>Siloxerus multiflorus</i>	Small wrinklewort
	E		<i>Stackhousia gummii</i>	Gunn's migronette
	V		<i>Triptilodiscus pygmaeus</i>	Common sunray
	V		<i>Velleia paradoxa</i>	Spur velleia
	R		<i>Villarsia exaltata</i>	Erect marsh flower
	R		<i>Viola cunninghamii</i>	Cunningham's violet
	R		<i>Vittadinia cuneata</i>	New Holland daisy
	R		<i>Vittadinia gracilis</i>	Graceful New Holland daisy
	R		<i>Vittadinia muelleri</i>	Narrow-leaf New Holland daisy
	R		<i>Xerochrysum bicolor</i>	White alpine everlasting
<b>Orchids</b>	R		<i>Caladenia caudata</i>	Tailed spider orchid
	E		<i>Corunastylis firthii</i>	Firth's midge orchid
	E		<i>Corunastylis morrisii</i>	Bearded midge orchid
	R		<i>Corunastylis nuda</i>	Tiny midge orchid
	E		<i>Diuris palustris</i>	Swamp diuris
	R		<i>Orthoceras strictum</i>	Horned orchid
	E		<i>Prasophyllum castaneum</i>	Chestnut leek orchid
	E		<i>Prasophyllum pulchellum</i>	Pretty leek orchid
	E		<i>Pterostylis wapstrarum</i>	Fleshy greenhood
	E		<i>Thelymitra antennifera</i>	Rabbit ears, lemon orchid
	E		<i>Thelymitra jonesii</i>	Sky-blue sun orchid
	E		<i>Thelymitra malvina</i>	Sun orchid
<b>Grasses</b>	R		<i>Agrostis diemenica</i>	Flat-leaf southern bent
	R		<i>Agrostis propinqua</i>	Alpine winter grass
	E		<i>Amphibromus macrorhinus</i>	Long nosed swamp wallaby grass
	R		<i>Amphibromus neesii</i>	Swamp wallaby grass
	E		<i>Austrodanthonia popinensis</i>	Roadside wallaby grass
	R		<i>Austrostipa bigeniculata</i>	Double-jointed speargrass
	R		<i>Austrostipa blackii</i>	Crested speargrass
	R		<i>Austrostipa nodosa</i>	Knotty speargrass
	R		<i>Austrostipa scabra</i>	Rough speargrass
	R		<i>Deyeuxia densa</i>	Heath bent grass
	R		<i>Lachnagrostis aequata</i>	Southern bent grass
	R		<i>Lachnagrostis billardierei</i> ssp. <i>tenuiseta</i>	Small awned blown grass
	R		<i>Lachnagrostis punicea</i> subsp. <i>filifolia</i>	Coast blown grass
	R		<i>Lachnagrostis punicea</i> subsp. <i>punicea</i>	Bristle blown grass
	R		<i>Lachnagrostis robusta</i>	Tall blown grass
	R		<i>Poa mollis</i>	Soft poa grass
	R		<i>Poa poiformis</i> var. <i>ramifer</i>	Island purple grass
	R		<i>Sporobolus virginicus</i>	Salt couch
<b>Other monocots</b>	R		<i>Aphelia pumilio</i>	Dwarf aphelia
	R		<i>Baumea articulata</i>	Jointed twig rush
	R		<i>Baumea gunnii</i>	Slender twig rush
	R		<i>Bolboschoenus caldwellii</i>	Sea club rush
	R		<i>Caesia calliantha</i>	Blue grass lily
	R		<i>Carex longibrachiata</i>	Drooping sedge
			<i>Carex tasmanica</i>	Curly sedge
	R		<i>Caustis pentandra</i>	Thick twist rush
		Prb	<i>Dianella amoena</i>	Matted flax lily
	R	Hs	<i>Dianella longifolia</i> var. <i>longifolia</i>	Pale or smooth flax lily
	R		<i>Hypoxis vaginata</i>	Sheathing yellow star
	R		<i>Isolepis habra</i>	Alpine club rush
	R		<i>Isolepis stellata</i>	Star club rush
	R		<i>Juncus amabilis</i>	Gentle juncus
	R		<i>Juncus prismatocarpus</i>	Branching rush
	R		<i>Juncus vaginatus</i>	Clustered rush
	R		<i>Lepilaena marina</i>	Sea water mat
	R		<i>Lepilaena preissii</i>	Slender water mat

**Forest Botany Manual: Module 4 – Freycinet Region**

	R		<i>Lepidosperma forsythii</i>	Stout rapier sedge
	R		<i>Lepidosperma tortuosum</i>	Twisting rapier sedge
	R		<i>Lepidosperma viscidum</i>	Sticky sword sedge
<b>Life form</b>	<b>Status</b>	<b>Pc</b>	<b>Botanical name</b>	<b>Common name</b>
<b>Other monocots</b>	R		<i>Potamogeton pectinatus</i>	Fennel pondweed
	R		<i>Ruppia tuberosa</i>	Tuberous tassel
	R		<i>Schoenus brevifolius</i>	Zig zag bog sedge
	V		<i>Tricoryne elatior</i>	Yellow rush lily
	R		<i>Tricostularia pauciflora</i>	Needlebog sedge
	V	Hs	<i>Xanthorrhoea arenaria</i>	Sand grasstree
		Hs	<i>Xanthorrhoea australis</i>	Austral grasstree
<b>Non-vascular</b>	R		<i>Hypotrachyna immaculata</i>	Lichen
	V		<i>Melanelia piliferella</i>	Lichen
	R		<i>Xanthoparmelia microphyllizan</i>	Lichen
	E		<i>Xanthoparmelia molliuscula</i>	Lichen
	R		<i>Xanthoparmelia oleosa</i>	Lichen

## Section 4 SITES OF POTENTIAL SIGNIFICANCE FOR FLORA

Information in this section is oriented towards sites rather than communities or species. It concentrates on environments within the region that have the potential to be directly or indirectly affected by forestry operations. This provides another approach to conservation of flora, which could be particularly useful for gaining an overview of potential botanical issues or values in an area. It could also be useful when there are problems with identifying species or communities in an area of proposed operations. However, it should not be used as a substitute for determining communities (Section 2) and priority species (Section 3) in an FPP area.

Species and communities of high conservation significance are often associated with particular environments. Sites can be identified by features of the physical environment (e.g. geology, altitude, landform) or the vegetation (e.g. dominant eucalypts, P.I. type). Local knowledge, coupled with use of aerial photographs and topographic, geology and P.I. maps, will help identify sites of potential significance. Many of these sites are of low or marginal timber value, and are not suitable for forestry activities or agricultural development. Such sites include dry ridgelines and river gorges. Other sites may be of greater commercial interest, for example forests on basalt and patches of relict rainforest. Many of the sites of potential significance in Freycinet Region contain species that are susceptible to *Phytophthora cinnamomi*.

Table 4A and 4B indicate forest and woodland sites that have the greatest potential to contain significant flora values.

**Table 4A: Sites that are often associated with priority communities or species.**

**Contact FPA in all cases if proposed operations could affect these sites.** The botanical significance of the site can then be evaluated from available information, or a vegetation survey may be needed. Areas of remnant forest that are proposed for conversion are included in this table, though they may not always contain priority communities or species.

**Table 4B: Sites that are occasionally associated with priority communities or species.**

**Assess these environments carefully. Contact FPA if priority species or communities are found in areas that could be affected by proposed operations.**

### Notes:

1. Sites supporting native non-forest vegetation types (e.g. heath, wetland, native grassland) have not been included in the table, though they may co-occur with forests and woodlands on many of the sites listed. Such vegetation often contains rare species, particularly in areas of the region where little non-forest vegetation remains. Contact FPA if areas of native non-forest vegetation could be affected by the proposed operation.
2. Management of remnant forests and woodlands, which often have important flora values, is treated in Section 6.

**Section 4 SITES OF POTENTIAL SIGNIFICANCE FOR FLORA**

The sites listed in this section are associated with species or communities with a priority for conservation in Freycinet Region. Information in the tables is not a substitute for information given in Section 2 and Section 3 of this module. Botanical advice should be sought for all sites with native non-forest vegetation.

**Table 4A: Sites often associated with priority communities or species. Contact FPA in all cases if these sites could be affected by operations.**

Site of potential significance	Main dominants	Main understorey	Main PI type	Reason for significance	Example locations
<b>Forests and woodlands on Tertiary ironstone, gravels or Recent sand or alluvium</b> in non-coastal areas	<i>E. amygdalina</i> , <i>E. viminalis</i> , <i>E. pauciflora</i> , <i>E. ovata</i>	Heathy, sedgy, or grassy	E4, E5	Priority communities and species (e.g. <i>Stenanthemum pimeleoides</i> )	Cranbrook, Fingal Valley
<b>River flats, frost hollows and marshes</b>	<i>E. pauciflora</i> , <i>E. viminalis</i> , <i>E. ovata</i> , tea-tree, paperbark	Variable – often scrubby, sedgy or grassy	E3, E4, E5, T, S, Vz or Wg	Priority communities and species (e.g. <i>Acacia axillaris</i> , <i>Euphrasia scabra</i> , <i>Isoetes drummondii</i> )	Thompsons Marsh, Dukes Marsh, Benjafields Marsh, Wrights Marsh, Flagstaff Marsh, Wet Gun Swamp
<b>Dry ridges, knolls, and upper slopes</b> often with high rock cover (mainly dolerite, also granite or sandstone)	<i>E. amygdalina</i> , <i>E. pulchella</i> , <i>E. viminalis</i> , <i>E. globulus</i> , she-oak	Grassy, shrubby or almost bare	E4 or E5; S or Wr often present	Priority communities and vascular and non-vascular species (e.g. <i>Spyridium lawrencei</i> , <i>Ozothamnus lycopodioides</i> , <i>Eucalyptus barberi</i> )	Mt Andrew, Cherry Tree Hill, Ringrove Razorback, The Thumbs, The Hazards
<b>Rocky outcrops</b> including plates, large boulders, cliffs and scree fields	Variable	Grassy, shrubby or bare	E4 or E5; S or Wr	Localised vascular and non-vascular species; susceptibility to disturbance	Nichols Cap, Mt Amos
<b>Vegetation with grasstrees</b> ( <i>Xanthorrhoea</i> species)	<i>E. amygdalina</i> , <i>E. pulchella</i> ,	Variable – often heathy	E4, E5: S	Priority species and high susceptibility to <i>Phytophthora</i>	Mt Allen, Freycinet NP
<b>Rocky gorges and creeklines</b> particularly with N or W orientation or aspect	<i>E. pulchella</i> , she-oak, Oyster Bay pine	Variable, often scrubby or shrubby	Variable	Priority communities and species (e.g. <i>Callitris oblonga</i> , <i>Epacris apsleyensis</i> , <i>Acacia axillaris</i> , <i>Bertya tasmanica</i> )	Paradise Gorge, Hardings Falls, Bluff River Gorge
<b>Recent sands in coastal areas</b>	<i>E. globulus</i> , <i>E. viminalis</i>	Variable, often scrubby	E4, E5	Priority communities	Rheban Spit, McRaes Isthmus (Maria Island), Dolphin Sands
<b>Exposed coastal sites</b> , especially on dolerite	She-oak, Oyster Bay pine, often sparse	Coastal shrubs	E5, S, Wr	Priority communities and species (e.g. <i>Allocasuarina crassa</i> , <i>Euphrasia phragmostoma</i> , <i>Euphrasia amphisysepala</i> )	Cape Huay, Cape Bernier

**Table 4B: Sites occasionally associated with priority communities or species. Assess these environments carefully. Contact FPA if priority communities or species could be affected by operations.**

Site of potential significance	Main dominants	Main understorey	Main PI type	Reason for significance	Example locations
<b>Very humid watercourses and slopes</b> generally with a S or E aspect, in fire shadow locations	Sassafras, myrtle, ( <i>E. regnans</i> , <i>E. obliqua</i> , <i>E. delegatensis</i> )	Rainforest	S or T; >E3 often present	Relict rainforest and other priority communities. Priority species (e.g. <i>Cyathea cunninghamii</i> , <i>Phyllocladus aspleniifolius</i> )	Big Sassy Creek, upper Buxton River, Little Beach Creek
<b>Coastal cloud-trapping landforms</b>	Sassafras, myrtle, celery-top pine, <i>E. urnigera</i>	Rainforest	S or T, >E3 often present	Relict rainforest and other priority communities. Priority species (e.g. <i>Phyllocladus aspleniifolius</i> )	Macgregor Peak, Mt Walter, Tatnells Hill area
<b>High altitude peaks and plateaux</b> mainly above 800 m	<i>E. delegatensis</i> , <i>E. coccifera</i>	Shrubby dry sclerophyll	<E4	Priority communities and species (e.g. <i>Eucalyptus coccifera</i> , <i>Eucalyptus gunnii</i> , <i>Eucalyptus archeri</i> )	Moaners Tier, Snow Hill, Middle Peak
<b>Dolerite-sediment contact zones on Central East Coast</b> where sedimentary substrate has thin overburden of dolerite talus	<i>E. amygdalina</i> , <i>E. tenuiramis</i> , <i>E. viminalis</i> , Oyster Bay Pine	Shrubby dry sclerophyll	E4	Priority communities and species (e.g. <i>Epacris limbata</i> )	Apsley River area
<b>Grassy forests, woodlands and other ecosystems</b>	<i>E. viminalis</i> , <i>E. globulus</i> , <i>E. ovata</i> , <i>E. pauciflora</i>	Grassy	E4, E5, Vz	Priority communities and species (e.g. <i>Austrostipa scabra</i> , <i>Velleia paradoxa</i> )	Green Tier Creek

## Section 5 EVALUATION OF OTHER FLORA ISSUES

Section 5 covers issues that need to be considered by FPOs to ensure that operations comply with the *Forest Practices Code* and other current legislation and policies. Issues discussed in this section will not be relevant to all FPP areas.

Some topics have already been covered to some extent in previous sections of the module (for example, Section 2 identifies forest communities that may need prescriptions related to *Phytophthora cinnamomi*). However, they are also treated in Section 5, because they are dealt with separately in the FPP *Flora Evaluation Sheet*. Reference to *Flora Technical Notes* may be needed.

Additional topics may be introduced to this section (and the *Flora Evaluation Sheet*) as information becomes available from research, and if there are changes to legislation, policies and codes of practice.

### WEED AND DISEASE MANAGEMENT

Flora values in many forest and scrub communities can be adversely affected by the introduction of disease and exotic plants. Section D3.1 of the *Forest Practices Code* gives guidelines to reduce the risk of weeds and disease being introduced through forestry operations. Quarrying, roading and road use are generally of more concern than logging and regeneration activities.

Diseases and weeds that can seriously threaten flora (and fauna) values are discussed below. Many other weeds and disease can affect wood production and plant species and communities – some of these are detailed in information available from Forestry Tasmania.

#### *Phytophthora cinnamomi*

*Phytophthora cinnamomi* (often called root rot fungus) is a disease that attacks the roots of many native species. Some plants die rapidly (e.g. banksias, grasstrees), while others (e.g. several eucalypt species) only show signs of disease in periods of drought or other stress. Many threatened species are highly susceptible. Open vegetation in relatively moist, lowland environments, such as dry sclerophyll forest, scrub, heath and moorland, are most at risk from *Phytophthora*. The resultant reduction in plant diversity and resources (e.g. nectar, pollen and shelter) has flow-on effects to fauna. Information on *Phytophthora* is given in *Flora Technical Note 8*.

*Phytophthora* is widespread in lowland areas of Freycinet Region, across all land tenures. Cool soil temperatures in wet forests and at higher altitudes tend to inhibit the disease, but opening up the canopy (e.g. by tracks) can cause local infestations.

*Phytophthora* has been introduced to many areas by soil carried on vehicles and machinery, but other sources include the boots of wandering people and the feet of wandering animals. Once established, it is impossible to eradicate, and can spread rapidly in surface run-off and groundwater percolation. The risk of spreading *Phytophthora* can be reduced by machinery hygiene, use of *Phytophthora*-free material in road construction, and attention to infrastructure planning (e.g. roads located on ridgelines will place a larger area of susceptible vegetation at risk than roads located on lower slopes).

Over sixty *Phytophthora* Management Areas, containing species or communities that are particularly susceptible to the pathogen, have been delineated – mainly on public land in lowland areas of the State. About a quarter of these are located in Freycinet Region (many in existing reserves).

Location of *Phytophthora* Management Areas and records of *Phytophthora* are given in databases that FPOs need to use to complete the FPP *Flora Evaluation Sheet* (Module 1 details how to access these databases). Forest communities that are susceptible to *Phytophthora* are indicated in *Flora Technical Note 8*. They are also indicated (#) in the community tables in Section 2 of this module.

FPOs need to refer to *Flora Technical Note 8* if:

- *Phytophthora* has been recorded from the proposed operational area;
- the operation will affect a highly susceptible forest community; or
- the operation is located within a *Phytophthora* Management Area.

Specialist advice should be sought if prescriptions in *Flora Technical Note 8* cannot be applied. Non-forest vegetation that could be affected by *Phytophthora* should routinely be referred to FPA.

## Myrtle wilt

Myrtle wilt is a disease of myrtle (*Nothofagus cunninghamii*) caused by the fungus *Chalara australis*. It kills infected trees, and can spread to other trees by root contact. Myrtle wilt occurs naturally in undisturbed forest. Damage to stems increases the chance of infection. *Flora Technical Note 9* provides guidelines to reduce the risk of infection to retained myrtles in operational areas. Myrtle has a very restricted occurrence in Freycinet Region, with almost all sites being in reserves (e.g. Douglas-Apsley NP, Tasman NP). Operations with the potential to adversely affect rainforest in formal reserves or Special Management Zones (Flora) should be routinely referred to FPA.

## Threatening weed species

Weed species can colonise disturbed sites associated with forestry operations, particularly when operations are close to agricultural land. Some weed species (e.g. thistles) decrease in abundance as understoreys re-establish in the regenerating forest. Other weed species are more persistent in forest – they include species with seeds that remain viable for a long time (e.g. gorse and broom), and species that are capable of vegetative regeneration (e.g. blackberry). Open sites, such as road verges, tracks, landings, quarries and bridge approaches, provide good environments for weeds to establish and persist. Weeds can also take advantage of disturbance (including burns associated with forest management) to establish in areas of non-forest vegetation (e.g. moorlands and native grasslands). Any infestation provides a launching pad, which allows the weed species to colonise other sites in the general area – through seeds dispersed by wind, birds, water movement, livestock or other land use (including road construction and use, and forestry operations).

There are legislative requirements under the Tasmanian *Weed Management Act* for land managers to control declared weed species. The required course of action will depend on the circumstances, including characteristics of the species, and extent of infestation at the site and in the municipality. The DPIWE website ([www.dpiwe.tas.gov.au](http://www.dpiwe.tas.gov.au)) gives details of the Act and Statutory Weed Management Plans for declared weed species. Declared weed species of most concern in forest environments include gorse, English broom, Spanish heath, ragwort, blackberry and pampas grass. A greater range of declared weed species may be present on plantation sites, or areas of previously cleared land proposed for plantation establishment. Control through hygiene and active management is particularly important where threatening exotics have the potential to spread into reserves or other areas of native forest where they are rare or absent.

## REMNANT FOREST AND WOODLAND

The *Forest Practices Code* supports the maintenance of remnant forest in those parts of the state where native vegetation has been extensively cleared. The RFA also requires that the values of remnant vegetation are considered at a regional level as a part of forest practices planning.

For the purpose of assessing FPP flora values, remnant forests and woodlands comprise stands that are:

- greater than 1 ha in area, and
- separated by more than 2 km from the closest area of native forest or woodland that exceeds 20 ha in area.

In some cases, remnant forests in Freycinet Region contain communities (e.g. *E. ovata* forest) and species that have a high priority for conservation. Other remnants contain communities that are more widespread and better reserved. Remnant vegetation will differ greatly in its condition – from sites with an understorey dominated by native species, to sites with understoreys having a high proportion of exotic shrubs or pasture grasses. In parts of the region with a long history of modification from agriculture and plantation forestry, such as parts of the Eastern Midlands, remnant vegetation may be scattered and degraded by a combination of land use, edge effects and attrition of species over the years. However, even substantially disturbed remnants can play an important role in maintaining flora and fauna at a local to subregional scale.

In all cases, remnants in FPP areas need to be carefully evaluated. Those containing communities (Section 2), species (Section 3) or sites of potential significance (Section 4) need to be referred to FPA for specialist advice. In addition, FPA needs to be contacted for any proposed operation (typically clearance for plantation or agriculture) involving conversion of remnant vegetation. Advice concerning the operation will take account of the composition, extent, condition and context of the remnant.

## MANAGEMENT OF FORESTRY OPERATIONS IN VICINITY OF RESERVES

This section deals with forestry operations (logging, roading, quarrying, plantation establishment, regeneration treatments) that are within or adjacent to formal reserves. Formal reserves are gazetted reserves on public land (e.g. National Parks, State Reserves, Forest Reserves) and reserves on private land that have been registered on property titles (e.g. reserves established through the Private Forest Reserves Program). This section also deals with operations that could affect areas categorised by Forestry Tasmania as Special Management Zones (Flora). Comments may also be relevant for other areas being managed for conservation on public and private land.

Botanical values in reserves adjacent to proposed operational areas will often be identified in the assessments of plant communities (Section 2); priority species (Section 3) and sites of potential significance (Section 4).

Some reserves incorporate buffers or have management systems that are designed to prevent, absorb or reduce disturbance associated with adjacent or nearby land use. However, good forest practices planning needs to take account of potentially adverse effects on botanical values (and other values) in all reserves. Such effects could include:

- introduction or increased incidence of weeds (including wildlings of pines or non-indigenous eucalypts sown or planted in the FPP area);
- introduction or increased incidence of disease (*Phytophthora* and myrtle wilt are of most concern – see above);
- change in microclimate in reserve [this is of most concern when vegetation along the reserve boundary contains localised wet forest vegetation (e.g. rainforest) which is susceptible to warmer and drier conditions];
- increased risk of fire entering the reserve (particularly when vegetation in the reserve is upslope of the forestry operation, and contains fire-sensitive plant communities or species).

The effect of forestry operations will depend on:

- attributes of the FPP area;
- attributes of the adjacent reserve and its vegetation (plant species and communities);
- the type of operation;
- regeneration treatment or post-operational land use.

FPOs need to consider carefully all these factors. Advice should be sought from FPA if the vegetation in the reserve has the potential to be adversely affected by an adjacent forestry operation and subsequent land use. FPA must be notified in all cases where a proposed operation is within a formal reserve or Special Management Zone (Flora).

## Section 6 ASSESSING THE FLORA VALUES OF AN AREA


The main aim of the *Forest Botany Manual* is to allow Forest Practices Officers, and others involved with preparing Forest Practices Plans, to comply with the requirements of the *Forest Practices Code* and other policies and legislation. A similar assessment process can also be used for other areas where botanical values need to be considered.

This section uses a flow diagram to summarise the actions FPOs should take after they have assessed the vegetation of an area. It is based on information contained in sections of the module dealing with forest communities (Section 2), priority species (Section 3), sites of potential significance for flora (Section 4) and other flora issues (Section 5). Relevant sections of the manual are indicated on the left side of the flow diagram. The FPP *Flora Evaluation Sheet* has a similar format to the diagram. An example of a completed *Flora Evaluation Sheet* is given in Module 1.

Note the following points:

1. The term 'FPP area' is used to describe the area under assessment – this may extend outside the proposed harvest area (e.g. cable tailhold areas).
2. The flora evaluation needs to determine if the operation will affect flora values in adjacent areas. Conversion of native vegetation has greater potential to affect nearby vegetation than native forest operations. The FPA needs to be contacted for any FPP where conversion of native vegetation is proposed next to gazetted reserves (public or private) or Special Management Zones or other informal reserves established to protect flora values.
3. If the assessment indicates that FPA should be contacted:
  - Advice on botanical issues can be sought from the FPA Botany or Ecology programs. A discussion by phone or email may allow the botanical issues to be resolved, or clarified prior to more formal notification.
  - The normal notification process will need to be followed in most situations. Forward the FPP *Flora Evaluation Sheet* to the FPA Ecologist. Additional information that will be useful includes maps showing distribution of priority communities or priority species in the FPP area. Other information that may be needed includes past and proposed land use, site characteristics (e.g. geology) and other natural or cultural values in the area. Some of this information is required for the FPP cover sheet.
  - FPOs need to consider, and discuss with FPA if necessary, if values protected by reservation or prescription would be adversely affected by logging, regeneration practices or other activities related to the forestry operation (e.g. if regeneration burning in a coupe could affect a threatened species in an exclusion zone.)
4. **It is essential that relevant forest planners and workers are aware of agreed recommendations (e.g. exclusion of areas from coupes or roadline; procedures for control of weeds and diseases; fire management prescriptions).**

**Section 5 ASSESSING THE FLORA VALUES OF AN AREA**



Document Control Log Table

Document Summary Information

Document name	FPA document version control guidelines
Version	1.1
Trim record	2011/42573
Owner	Biodiversity Program
Author(s)	Fred Duncan
Release date	2005
Release Approved by	FPA
Release status	Final

Version Control

Version	Date	Author(s)	Summary of changes
0.1	2 May 2011	Tim Leaman	Document creation in TRIM

Stages required for release outside FPA

Category of advice		B2
Stages	Required/not required	Completed (date)
Specialist	Required	2005
Line Manager	Required	2005
Peer/stakeholder review (FPA staff)	Required	2005
CFPO	Required	2005
FPAC	Not required	
Board	Not required	